

# **CHILDRENS PLUS COUGH AND RUNNY NOSE- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide suspension Walgreens**

-----  
**Drug Facts**

## **Active ingredients (in each 5 mL)**

### **Acetaminophen 160 mg**

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

## **Purposes**

### **Pain reliever/fever reducer**

Antihistamine

Cough suppressant

## **Uses**

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - cough
  - headache
  - sore throat
  - sneezing and runny nose
- temporarily reduces fever

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL each) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

### **Ask a doctor before use if the child has**

- Liver disease
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

### **Ask a doctor or pharmacist before use if the child is taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dose (see overdose warning )**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain or cough gets worse or lasts for more than 5 days
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### **Directions**

- **this product does not contain directions or complete warnings for adult use.**
- mL = milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

### Other Information

- store between 20-25°C (68-77°F). Do not refrigerate

### Inactive ingredients

acesulfame potassium, alcohol, carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose purified water, sodium benzoate, sorbitol, xanthan gum

### Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### Principal Display Panel

Compare to Children's Tylenol® Plus Cough & Runny Nose active ingredients††

#### Children's

#### Cough & Runny Nose

ACETAMINOPHEN /

PAIN RELIEVER / FEVER REDUCER

CHLORPHENIRAMINE MALEATE /

ANTIHISTAMINE

DEXTROMETHORPHAN HBr /

COUGH SUPPRESSANT

- Relieves fever, sore throat, aches, sneezing, runny nose & cough
- Oral suspension

- ALCOHOL 0.005%

AGES 6 - 11 YEARS

FL OZ (mL)

CHERRY FLAVOR

DOSAGE CUP PROVIDED

††This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Cough & Runny Nose.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

walgreens.com

**Package Label**

**Drug Facts (continued)**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

■ **this product does not contain directions or complete warnings for adult use**

- mL = milliliter ■ shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients:** acesulfame potassium, alcohol, carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, xanthan gum

**Questions or comments?**  
Call 1-877-753-3636 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**PARENTS:**  
Learn about teen medicine abuse  
www.StepMedicineAbuse.org

Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey

<sup>††</sup>This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Cough & Runny Nose.

PLD-K246D FC004160

*Walgreens*

Compare to Children's Tylenol® Plus Cough & Runny Nose active ingredients<sup>††</sup>

**children's**

# Cough & Runny Nose

**ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER  
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE  
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT**



- Relieves fever, sore throat, aches, sneezing, runny nose & cough
- Oral suspension
- ALCOHOL 0.005%

**AGES 6-11 YEARS**



**CHERRY FLAVOR**

4 FL OZ (118 mL)

**Drug Facts**

**Active ingredients (in each 5 mL)**

Active ingredients (in each 5 mL)	Purposes
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant

**Uses**

- temporarily relieves the following cold/flu symptoms
- minor aches and pains ■ sore throat ■ cough
- sneezing and runny nose ■ headache
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL each) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has**

- liver disease
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Ask a doctor or pharmacist before use if your child is taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

**When using this product**

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2017 Walgreen Co.  
ITEM 510120 W10067-XXXX-L



3 11917 10359 4

ORG0717-F2

Lot No:

Exp. Date:

## WALGREENS Children's Cough & Runny Nose Cherry Flavor

### CHILDRENS PLUS COUGH AND RUNNY NOSE

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide suspension

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0297
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH)	DEXTROMETHORPHAN	5 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)

HYDROBROMIDE

in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0297-04	1 in 1 BOX	07/31/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/31/2015	

**Labeler** - Walgreens (008965063)