### SAFE-GUARD- fenbendazole granule Merck Sharp & Dohme Corp.

-----

## safe-guard® (fenbendazole)

**Directions:** The daily dose for Safe-Guard<sup>®</sup> Canine is 50 mg/kg (22.7 mg/lb) of body weight. You should weigh your dog to make sure you are using the right dose; i.e. correct gram amount and appropriate number of pouches.

Your dog must be treated with the following dose for 3 days in a row.

Dog Weight (lbs)	Pouch Size (Daily Dose)
10	1 gram
11 to 20	2 gram
21 to 30*	1 gram + 2 gram
31 to 40	4 gram
41 to 50*	1 gram + 4 gram
51 to 60*	2 gram + 4 gram
61 to 80*	Two 4 gram
Over 80*	
Mix daily dose with a small amount of the usual food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing.  Repeat the dose for 3 days in a row.	2000 CO

<sup>\*</sup> Combine pouches to obtain recommended daily dosage.

# **Recommended Deworming Schedule:**

Schedule may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as a general guideline.

- Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age.
- Treat the dam at the same time as the puppies.
- Dogs over 6 months of age should be dewormed at least twice each year.

**Diagnosis of Parasites:** Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard<sup>®</sup> Canine only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard<sup>®</sup> Canine, consult your veterinarian.

Roundworms and hookworms may pose serious human health risks; regular deworming

may help reduce those risks.

**Product Use:** Safe-Guard<sup>®</sup> Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

**Adverse Reactions:** In U.S. clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with the use of the product.

**Storage:** Store at controlled room temperature 68-77°F (20-25°C).

#### **Questions? Comments?**

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call **1-800-441-8272**.

#### **MERCK**

**Animal Health** 

Distributed by: Intervet Inc (d/b/a Merck Animal Health) Madison, NJ 07940

Rev. 07/22

Approved by FDA under NADA # 121-473

Fenbendazole (active ingred.) made in: see imprint. Formulated in Austria.

www.Safe-Guard.com

#### PRINCIPAL DISPLAY PANEL - 3 Pouch Carton - 10 lbs

**MERCK** 

**Animal Health** 

safe-guard® (fenbendazole)

Canine

Dewormer

For dogs only, 6 weeks and older.

Drug Facts:


Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

Uses: For the treatment and control of:

- Tapeworms (*Taenia pisiformis*)
- Roundworms (Toxocara canis, Toxascaris leonina)
- Hookworms
   (Ancylostoma caninum, Uncinaria stenocephala)
- Whipworms (*Trichuris vulpis*)

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

HUMAN WARNING: KEEP OUT OF REACH OF CHILDREN.



#### PRINCIPAL DISPLAY PANEL - 3 Pouch Carton - 20 lbs

MERCK Animal Health

safe-guard® (fenbendazole)

Canine

Dewormer

For dogs only, 6 weeks and older.

# Drug Facts:


Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

Uses: For the treatment and control of:

- Tapeworms (*Taenia pisiformis*)
- Roundworms
   (Toxocara canis, Toxascaris leonina)
- Hookworms
   (Ancylostoma caninum, Uncinaria stenocephala)
- Whipworms (*Trichuris vulpis*)

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

HUMAN WARNING: KEEP OUT OF REACH OF CHILDREN.



MERCK Animal Health

safe-guard® (fenbendazole)

Canine Dewormer

For dogs only, 6 weeks and older.

Drug Facts:

Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

Uses: For the treatment and control of:

- Tapeworms (*Taenia pisiformis*)
- Roundworms (Toxocara canis, Toxascaris leonina)
- Hookworms
   (Ancylostoma caninum, Uncinaria stenocephala)
- Whipworms (*Trichuris vulpis*)

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

HUMAN WARNING: KEEP OUT OF REACH OF CHILDREN.



Diagnosis of Parasites: Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard® Canine only kills Taenia species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard® Canine, consult your veterinarian.

Net Weight: 12 grams

Roundworms and hookworms may pose serious human health risks; regular deworming may help reduce those risks.

Product Use: Safe-Guard® Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

Adverse Reactions: In U.S. clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with the use of the product.

Storage: Store at controlled room temperature 68-77°F (20-25°C).

#### Questions? Comments?

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call 1-800-441-8272.

1-800-441-8272.

Distributed by:
Intervet Inc (d/b/a Merck Animal Health)
Madison, NJ 07940

Approved by FDA under NADA # 121-473
Fenbendazole (active ingred, made in: see imprint. Formulated in Austria.



 Dogs over 6 months of age should be dewormed at least twice each year.

 Colorfree

■ Treat the dam at the same time as the

Recommended Deworming Schedule:

general guideline.

Schedule may vary depending on the climate where you live and the activity of your dog.


The following schedule should be used as a

■ Weaned pups (6-8 weeks of age) should be

dewormed at 6, 8, 10 and 12 weeks of age.



colorfree 25mmx12mm



MERCK

#### SAFE-GUARD

fenbendazole granule

# Product Information Product Type OTC ANIMAL DRUG Item Code (Source) NDC:57926-483 Route of Administration ORAL

# Active Ingredient/Active Moiety Ingredient Name Basis of Strength FENBENDAZOLE (UNII: 621BVT9M36) (FENBENDAZOLE - UNII:621BVT9M36) FENBENDAZOLE (222 mg in 1 g

P	Packaging							
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date				
1	NDC:57926-483-01	3 in 1 CARTON						
1		1 g in 1 POUCH						

2	NDC:57926-483-02	3 in 1 CARTON	
2		2 g in 1 POUCH	
3	NDC:57926-483-04	3 in 1 CARTON	
3		4 g in 1 POUCH	

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
NADA	NADA121473	01/05/1979				

Labeler - Merck Sharp & Dohme Corp. (001317601)

Revised: 11/2024 Merck Sharp & Dohme Corp.