

BERKLEY AND JENSEN PAIN RELIEF EXTRA STRENGTH- acetaminophen tablet BJWC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BJWC Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain one or more of these ingredients

Questions or comments?

1-800-934-1204

Principal Display Panel

Compare to the active ingredient in Extra Strength Tylenol[®]

SEE NEW WARNINGS

EXTRA STRENGTH

PAIN RELIEF

ACETAMINOPHEN

PAIN RELIEVER / FEVER REDUCER

FOR ADULTS

GLUTEN FREE

500 CAPLETS

500 mg EACH

ACTUAL SIZE

Compare to the active ingredient in Extra Strength Tylenol[®] ** SEE NEW WARNINGS

berkley jensen

EXTRA STRENGTH
PAIN RELIEF ACETAMINOPHEN
PAIN RELIEVER/FEVER REDUCER
FOR ADULTS

GLUTEN FREE
gf

500 CAPLETS | 500 mg EACH ACTUAL SIZE

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - muscular aches
 - temporarily reduces fever
- headache
- backache
- boothache
- minor pain of arthritis
- premenstrual and menstrual cramps

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- redness
- itching
- swelling
- skin peeling
- skin blisters
- skin sores

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- fever lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- take 2 caplets every 6 hours while symptoms last
- do not take more than 10 caplets in 24 hours unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- ask a doctor

children under 12 years

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients camellia wax, corn starch, croscarmellose sodium, hydroxypropylcellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid *may contain one or more of these ingredients

Questions or comments? 1-800-934-1204

*This product is not manufactured or distributed by McKell Consumer Healthcare, distributor of Extra Strength Tylenol[®].

Every Berkley Jensen product comes with our **100% Money-Back Guarantee**. It's just one of the many ways a BJ's membership gives you MORE.

Distributed by:
BJ's Wholesale Club
25 Research Drive
Parsippany, NJ 07054
Questions? 1-800-934-1204

Manf. #075798098
8 88670 101803

: 46490 D3 F8

BERKLEY AND JENSEN PAIN RELIEF EXTRA STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:6839 1-484
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE (caplet)	Size	16mm
Flavor		Imprint Code	L484
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-484-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/18/2006	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/18/2006	

Labeler - BJWC (159082692)