PRIMO TOPICAL ANESTHETIC- benzocaine gel Tri State Dental

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use. Use only as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Keep out of reach of children.

Other Information

Store at 59°-86°F (15°-30°C). Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

1-800-232-6874 or www.primodentalproducts.com

99291551 Rev 11/2018

PEEL BACK to review drug facts

845 Third Avenue New York, NY 10022 800-232-6874 www.primodentalproducts.com

When using this product



Gluten Free

Net Content: 1 oz (30 g)

mouth symptoms do not improve if swelling, rash or fever develops, directed by a physician. If sore in 7 days; irritation, pain, or

 in children under 2 years of age edness persists or worsens; or see your physician promptly. for teething

Do not exceed recommended contact occurs, flush with water. Avoid contact with eyes. If

> may cause methemoglobinemia, a serious condition that must be

warning: Use of this product

Methemoglobinemia

Poison Control Center right away. dosage. If more than used for pain is accidentally swallowed, get medical help or contact a

If pregnant or breast feeding, Keep out of reach of children. ask a physician before use.

Directions

attention if you or a child in your

pale, gray or blue colored skin

care develops:

occur even if you have used this

product before. Stop use and

seek immediate medical

reduces the amount of oxygen

treated promptly because it

carried in the blood. This can

 Apply only amount needed to children under 2 years of age: the oral mucosa to prevent or relieve pain.

do not use

Store at 59° - 86°F (15 - 30°C). Other information

dizziness or lightheadedness

 shortness of breath rapid heart rate

Allergy Alert: Do not use on

fatigue or lack of energy

allergies to local anesthetics

patients with a history of

such as procaine, butacaine,

benzocaine or other "caine"

contain blue #1, green #3, green sodium saccharin, water. May flavoring, PEG 3350, PEG 400, **Inactive ingredients**

1-800-232-6874 or www.primodentalproducts.com Questions or comments? as a color additive.

headache

(cyanosis)

Protect from freezing.

yellow #5, (tartrazine), yellow #6, #5, red #3, red #28, red #40,

> Do not use anesthetics.

for more than 7 days unless

Benzocaine 200 mg (in each g) Active ingredients **Drug Facts** Purpose Use

Oral Anesthetic

For oral mucosal use only, as temporary relief of pain due to minor dental procedures. directed by dentist. For the Warnings

PRIMO TOPICAL ANESTHETIC

benzocaine gel

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69509-009
Route of Administration	DENTAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients	
Ingredient Name	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C YELLOW NO. 5 (UNII: 1753WB2F1M)	

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics			
Color	yellow	Score	
Shape		Size	
Flavor	BANANA	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69509-009- 30	30 g in 1 JAR; Type 0: Not a Combination Product	11/01/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	03/01/2014	

Labeler - Tri State Dental (176048478)

Revised: 1/2022 Tri State Dental