

ACETAMINOPHEN- acetaminophen tablet
LEADER/ Cardinal Health 110, Inc

8HR Muscle Aches & pain Relief
Acetaminophen Extended-Release Tablets USP, 650mg
Pain reliever/Fever Reducer

Active ingredient (in each caplet)

Acetaminophen USP, 650mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- muscular aches
- backache
- minor pain of arthritis
- toothache
- premenstrual and menstrual cramps
- headache
- the common cold
- temporarily reduces fever

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- More than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years of age and over

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

children under 12 years

- do not use

Other information

- store between 20-25°C (68-77°F)
- **do not use if foil inner seal is broken or missing**

Inactive Ingredients

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?

Contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.

PDP

end user :CRANULSS
 item :CVS APAP RR 100CT
 code # :
 size :2+3/32 x 2+3/32 x 4+3/16
 ref # :PP180604A view :2
 material :.016 SRS date :0



Important: Read product information before using. Keep the carton for important information.

Do not use if foil inner seal is broken or missing.

Drug Facts
Active ingredient (in each caplet) Purpose
 Acetaminophen USP 650 mg... Pain reliever/fever reducer

Uses
 temporarily relieves minor aches and pains due to:
 ■ muscular aches
 ■ headache
 ■ minor pain of arthritis
 ■ toothache
 ■ premenstrual and menstrual cramps
 ■ headache
 ■ the common cold
 ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount.
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening
 ■ hives
 ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
 Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Drug Facts (continued)
Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.




Directions
 ■ do not take more than directed (see overdose warning)
 adults and children 12 years of age and over
 ■ take 2 caplets every 8 hours with water
 ■ swallow whole; do not crush, chew, split or dissolve
 ■ do not take more than 6 caplets in 24 hours
 ■ do not use for more than 10 days unless directed by a doctor
 children under 12 years
 ■ do not use

Other information
 ■ store between 20-25°C (68-77°F)
 ■ do not use if foil inner seal is broken or missing

Inactive ingredients
 hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?
 contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 9:00 PM PST

Contains No Aspirin
 *This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® 8 HR Muscle Aches & Pain.



ACETAMINOPHEN acetaminophen tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0626
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg
Inactive Ingredients			

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYDROXYETHYL CELLULOSE (140 CPS AT 5%) (UNII: 8136Y38GY5)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	

Product Characteristics

Color	white (WHITE TO OFF WHITE COLORED)	Score	no score
Shape	OVAL (CAPSULE SHAPED, BICONVEX INTACT FILM COATED TABLETS)	Size	19mm
Flavor		Imprint Code	G;650
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0626-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/02/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	03/02/2023	

Labeler - LEADER/ Cardinal Health 110, Inc (063997360)

Revised: 12/2025

LEADER/ Cardinal Health 110, Inc