

**MEDIQUE MEDIPROXEN- naproxen sodium tablet, film coated**  
**Unifirst First Aid Corporation**

-----  
**Medique Mediproxen**

***Drug Facts***

***Active ingredient (in each tablet)***

Naproxen Sodium 220 mg

(Naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves minor aches and pains due to

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold

temporarily reduces fever

***Warnings***

**Allergy Alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning:**

NSAID, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you ever have had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially

important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

**Adults: (12 years of age and older)**

- take 1 tablet every 8 to 12 hours while symptoms last.
- for the first dose you may take 2 tablets within the first hour.
- do not exceed 2 tablets in any 8 to 12 hour period
- do not exceed 3 tablets in a 24 hour period

**Children under 12 years:** ask a doctor

***Other information***

- each tablet contains: sodium 20 mg
- store at 68-77°F (20-25°C)
- avoid high humidity and excessive heat above 104°F (40°C)
- tamper evident sealed packets
- do not use any open or torn packets

***Inactive ingredients***

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

***Questions or comments?*** 1-800-634-7680

Medique®

Mediproxen

Long-Lasting Pain Relief

This Package is for Households without Small Children

Pull to Open

Pain, Fever • Naproxen Sodium USP, 220 mg (NSAID)

50 Tablets

(50 x 1)


Tamper Evident Unit Dose Packets



**Medique**  
**Mediproxen**  
Long-Lasting Pain Relief

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

✓ Pain, Fever • Naproxen Sodium USP, 220 mg (NSAID)




**Medique**  
**Mediproxen**  
Long-Lasting Pain Relief

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Pull to Open

✓ Pain, Fever • Naproxen Sodium USP, 220 mg (NSAID)

50 Tablets (50 x 1)



**Medique**  
**Mediproxen**  
Long-Lasting Pain Relief

2379580K  
Rev. 7/8/22  
Pg. 32

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Naproxen Sodium USP 220 mg ... Pain reliever/fever reducer (Naproxen 220 mg (NSAID))  
Nonsteroidal anti-inflammatory drug

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ minor pain of arthritis ■ muscle aches ■ backache  
■ menstrual cramps ■ headache ■ toothache  
■ the common cold ■ temporarily reduces fever

**Warnings**  
**Alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin redness ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 65 or older  
■ have stomach ulcers or bleeding problems  
■ take a blood thinner (anticoagulant) or steroid drug

**Drug Facts (continued)**

**When using this product:**  
■ use with food or milk if stomach upset occurs

**Stop use and ask a doctor if:**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing  
■ weakness in one part or side of body  
■ slurred speech ■ leg swelling  
■ pain gets worse or lasts for more than 10 days  
■ ear pain worse or lasts for more than 3 days  
■ you have difficulty swallowing  
■ it feels like the pill is stuck in your throat  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**

**Directions**  
■ do not take more than directed  
■ the smallest effective dose should be used  
■ drink a full glass of water with each dose

**Adults:** (12 years of age and older)  
■ Take 1 tablet every 8 to 12 hours while symptoms last.  
■ For the first dose you may take 2 tablets within the first hour.  
■ do not exceed 2 tablets in any 8 to 12 hour period.  
■ do not exceed 3 tablets in a 24 hour period.

**Children under 12 years:** ask a doctor

**Other information**  
■ each tablet contains: sodium 20 mg  
■ store at 68°F (20°C).  
■ avoid high humidity and excessive heat above 104°F (40°C).  
■ tamper evident sealed packets.  
■ do not use any open or torn packets.

**Inactive ingredients**  
croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, polyethylene glycol, pregelatinized starch, tac, titanium dioxide

**Questions or comments?** 1-800-634-7880

Retain carton for complete product information      Tamper Evident Unit Dose Packets      Retain carton for complete product information

# MEDIQUE MEDIPROXEN

naproxen sodium tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-237
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	blue (blue)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	9mm

<b>Flavor</b>		<b>Imprint Code</b>	I;3	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-237-50	50 in 1 BOX	12/30/2008	
1		1 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-237-33	100 in 1 BOX	12/30/2008	
2	NDC:47682-237-46	1 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-237-46	1 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA079096	12/30/2008		

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 1/2024

Unifirst First Aid Corporation