

**COLD AND FLU RELIEF NIGHTTIME- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled  
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

**Pain reliever/Fever Reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do

not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

### **Other information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

Call **1-888-309-9030**

### **Principal Display Panel**

Compare to the active ingredients in Vicks® Nyquil® Cold & Flu LiquiCaps®\*\*

Night Time

Cold and Flu Relief

Acetaminophen- Pain Reliever/Fever Reducer

Dextromethorphan HBr- Cough Suppressant

Doxylamine Succinate- Antihistamine

Relieves:

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing

Alcohol Free

Softgel†

(†Liquid Filled Capsules)

\*\*This product is not manufactured or distributed by the Procter & Gamble Company.

Vicks®, NyQuil®, and LiquiCaps® are registered trademark of the Procter & Gamble Company

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**Package Label**

<p><b>Drug Facts</b></p> <p><b>Active Ingredients (in each softgel)</b></p> <p>Acetaminophen 325 mg  Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 15 mg  Cough suppressant</p> <p>Doxylamine succinate 6.25 mg  Antihistamine</p>	
<p><b>Uses</b></p> <p>temporarily relieves common cold and flu symptoms</p> <p>sore throat ■ headache ■ minor aches and pains</p> <p>fever ■ runny nose and sneezing</p> <p>cough due to minor throat and bronchial irritation</p>	
<p><b>Warnings</b></p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen</p> <p>3 or more alcoholic drinks every day while using this product</p> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
<p><b>Do not use</b></p> <p>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p>If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>	
<p><b>Directions</b></p> <p>do not take more than directed (see Overdose warning)</p> <p>do not take more than 4 doses in 24 hours</p>	
<p><b>Drug Facts (continued)</b></p> <p>Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema</p> <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> <li>taking the blood thinning drug warfarin</li> <li>taking sedatives or tranquilizers</li> </ul> <p>When using this product ■ do not exceed recommended dosage</p> <p>excitability may occur, especially in children</p> <p>avoid alcoholic drinks ■ marked drowsiness may occur</p> <p>alcohol, sedatives, and tranquilizers may increase drowsiness</p> <p>be careful when driving a motor vehicle or operating machinery</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>pain or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present ■ new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	

**DG™ health**

# Night Time Cold & Flu Relief



Acetaminophen - Pain Reliever/Fever Reducer  
Dextromethorphan HBr - Cough Suppressant  
Doxylamine Succinate - Antihistamine

**DG™ health**

Compare to the active ingredients of Vicks® NyQuit® Cold & Flu LiquiCaps®\*\*

# Night Time Cold & Flu Relief



Acetaminophen - Pain Reliever/Fever Reducer  
Dextromethorphan HBr - Cough Suppressant  
Doxylamine Succinate - Antihistamine

Relieves:

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing

Alcohol Free



**48 Softgels†**

(†Liqued Filled Capsules)

Actual Softgel Size

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.  
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

**PARENTS:**  
Learn more! Visit [www.StopMedicineForum.org](http://www.StopMedicineForum.org)

Lot No.:  
Exp. Date:



**100%**  
Satisfaction  
Guaranteed!  
(888) 309-9030

PLD-F585B F000284  
DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

**Other information**

- store between 15-30°C (59-86°F) ■ avoid excessive heat
- label to ensure correct dosing
- when using other Daytime or Nighttime products, carefully read each
- children under 12 years: do not use
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours

**Drug Facts (continued)**

**Drug Facts (continued)**

**inactive ingredients**: D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**  
Call 1-888-309-9030

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

**DOLLAR GENERAL HEALTH Nighttime Cold & Flu Relief**

**COLD AND FLU RELIEF NIGHTTIME**

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-599
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O921CV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1D156QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-599-48	48 in 1 CARTON	12/27/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/27/2019	

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 3/2024

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)