

**VICKS DAYQUIL NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION- acetaminophen, dextromethorphan hbr, diphenhydramine hcl, phenylephrine hcl
Procter & Gamble Manufactura S de RL de CV**

**Vicks® DayQuil™ HOT REMEDY Vicks® NyQuil™ HOT REMEDY
RAPID RELIEF
MULTI-SYMPTOM
COLD & FLU PLUS CONGESTION**

**Vicks® DayQuil™ HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD & FLU +
CONGESTION**

Drug Facts

DayQuil™

Active Ingredient (in each Packet)

Acetaminophen 500 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages

- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

**When using this product,
do not use more than directed**

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- **each packet contains:** potassium 10 mg

- phenylketonurics: contains phenylalanine 64 mg per dose
- do not exceed 25°C.

Inactive ingredients:

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

Questions?

1-800-362-1683

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

Vicks® NyQuil™ HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD & FLU + CONGESTION

Drug Facts

NyQuil™

Active Ingredient (in each Packet)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation

- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose
- sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease

- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **Do not use more than directed**
- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- **each packet contains:**potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- do not exceed 25°C

Inactive ingredients

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

Questions?

1-800-362-1683

Tamper Evident:Do not use if sachet is opened or torn

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 16 packet package

MAX STRENGTH

Vicks® DayQuil™ HOT REMEDY

RAPID RELIEF

MULTI-SYMPTOM

COLD & FLU + CONGESTION

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy

Vicks® NyQuil™ HOT REMEDY

RAPID RELIEF

MULTI-SYMPTOM

COLD & FLU + CONGESTION

Acetaminophen, Diphenhydramine HCl, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose, Cough

Nasal Congestion, Sinus Pressure

Nighttime Relief

HONEY LEMON FLAVORED

10 DAYQUIL POWDER PACKETS; 6 NYQUIL POWDER PACKETS; 16 TOTAL POWDER PACKETS

Part 1 of 2**VICKS NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION**

acetaminophen, diphenhydramine hcl, phenylephrine hcl powder, for solution

Product Information**Item Code (Source)** NDC:58933-551**Route of Administration** ORAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 5 g
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 g
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 5 g

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SUCROSE (UNII: C151H8M554)	
ASPARTAME (UNII: Z0H242BBR1)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor	HONEY (LEMON)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:58933-551-01	5 g in 1 PACKET; Type 0: Not a Combination Product	
---	------------------	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2022	

Part 2 of 2

VICKS DAYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPATOM COLD AND FLU PLUS CONGESTION

acetaminophen, dextromethorphan hbr, phenylephrine hcl powder, for solution

Product Information

Item Code (Source)	NDC:58933-550
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 g
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 5 g
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 g

Inactive Ingredients

Ingredient Name	Strength
SUCROSE (UNII: C151H8M554)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	
TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
ASPARTAME (UNII: Z0H242BBR1)	

Product Characteristics

Color	white	Score	
Shape		Size	

Flavor	HONEY (LEMON)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-550-01	5 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2022	06/30/2027

Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)

Revised: 1/2026

Procter & Gamble Manufactura S de RL de CV