

**ASPIRIN- aspirin tablet, chewable**  
**Advance Pharmaceutical Inc.**

-----  
**ASPIRIN 81mg**

**Active Ingredient**  
**(in each tablet)**

Aspirin 81 mg (NSAID) \*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain Reliever

**Use**

- temporarily relieves minor aches and pains
- for other uses, see your doctor, but do not use for more than 10 days without consulting your doctor because serious side effect may occur.

**Warnings**

**Reye's Syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early signs of Reye's syndrome, a rare but serious illness.

**Allergy Alert:**Aspirin may cause a severe allergic reaction which may include

- hives
- asthma (wheezing)
- shock
- facial swelling

**Stomach Bleeding Warning:**This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child

- has had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, others)
- takes more or for a longer time than directed

**Do not use**

- if the child has ever had an allergic reaction to any other pain reliever / fever reducer
- for at least 7 days after tonsillectomy or oral surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn

- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting and diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child is taking a diuretic
- child has asthma

**Ask a doctor or pharmacist before use if the child is**

- taking a prescription drug for diabetes, gout or arthritis
- taking any other drug
- under a doctor’s care for any serious condition

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- pain gets worse or lasts more than 5 days
- any new symptoms appear
- redness or swelling is present in the painful area
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- administer only under adult supervision
- if possible, use weight to dose, otherwise use age
- repeat dose every 4 hours while symptoms last
- do not give more than 5 doses in 24 hours
- drink water with each dose

<b>Weight (lbs.)</b>	<b>Age (years)</b>	<b>Dose</b>
77 - 83	11 to under 12	4 - 6 tablets
66 - 76	9 to under 11	4 - 5 tablets
46 - 65	6 to under 9	4 tablets
36 - 45	4 to under 6	3 tablets
32 - 35	3 to under 4	2 tablets
under 32	under 3	ask a doctor

**DRINK A FULL GLASS OF WATER WITH EACH DOSE**

- **adults and children 12 years and older** :take 4 to 8 tablets every 4 hours while symptoms last, but not more than 48 tablets in 24 hours.
- **Children under 12 years:**ask a doctor

### **Other Information**

- store at 15-30 °C (59-86 °F)

### **Inactive Ingredients**

corn starch, FD&C Yellow #6 alum. lake, magnesium stearate, orange flavor, silicon dioxide, sugar

### **Questions or Comments**


Call 631-981-4600, 8.30am – 4.30 pm ET Monday-Friday

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Manufactured by: Advance Pharmaceutical, Inc. Holtsville, NY 11742**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**DIE 0183**  
 1 5/8 x 1 5/8 x 3 1/8  
 FLAT SIZE 7.125X 7.46875



Exp. Date: \_\_\_\_\_  
 Lot No.: \_\_\_\_\_  
 9 95600 17714 0033 0  
 CA 1112

**Drug Facts (continued)**  
**Do not use**  
 ■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ for at least 7 days after tonsillectomy or oral surgery  
**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to your child  
 ■ child has a history of stomach problems, such as heartburn  
 ■ child has not been drinking fluids  
 ■ child has lost a lot of fluid due to vomiting or diarrhea  
 ■ child has high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ child is taking a diuretic  
 ■ child has asthma  
**Ask a doctor or pharmacist before use if the child is**  
 ■ taking a prescription drug for diabetes, gout or arthritis  
 ■ taking any other drug  
 ■ under a doctor's care for any serious condition  
**Stop use and ask a doctor if**  
 ■ child experiences any of the following signs of stomach bleeding:  
 ■ feels faint ■ vomits blood  
 ■ has bloody or black stools  
 ■ has stomach pain that does not get better  
 ■ pain gets worse or lasts more than 5 days  
 ■ any new symptoms appear  
 ■ redness or swelling is present in the painful area  
 ■ ringing in the ears or a loss of hearing occurs ▶

**Drug Facts (continued)**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
 ■ this product does not contain directions or complete warnings for adult use  
 ■ administer only under adult supervision  
 ■ if possible, use weight to dose, otherwise use age  
 ■ repeat dose every 4 hours while symptoms last  
 ■ do not give more than 5 doses in 24 hours  
 ■ drink water with each dose  


Weight (lbs.)	Age (years)	Dose
77 - 83	11 to under 12	4 - 6 tablets
66 - 76	9 to under 11	4 - 5 tablets
46 - 65	6 to under 9	4 tablets
36 - 45	4 to under 6	3 tablets
32 - 35	3 to under 4	2 tablets
under 32	under 3	ask a doctor

**Other information**  
 store at room temperature 15°-30°C (59°-86°F)  
**Inactive ingredients** corn starch, FD&C yellow #6 alum. lake, magnesium stearate, orange flavor, silicon dioxide, sugar  
**Questions or comments?**  
 call 631-981-4600, 8:30 am - 4:30 pm ET  
 Monday - Friday

**CHILDREN'S ASPIRIN (NSAID)**  
 Pain Reliever  
 1 1/4 gr (81 mg)  
 36 TABLETS  
 Chewable • Orange Flavor  
 Advance Pharmaceutical Inc.  
 Compare to Active Ingredient in St. Joseph® Chewables  
 See New Warnings Information  
 NDC 17714-009-36

**SAVE CARTON FOR COMPLETE DRUG FACTS**  
**Drug Facts**  
**Active ingredient (in each tablet)** Aspirin 81 mg (NSAID)  
**Purpose** Pain reliever  
 \*nonsteroidal anti-inflammatory drug  
**Use** temporarily relieves minor aches and pains  
**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include  
 ■ hives ■ asthma (wheezing)  
 ■ shock ■ facial swelling  
**Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if your child:  
 ■ has had stomach ulcers or bleeding problems  
 ■ takes a blood-thinning (anticoagulant) or steroid drug  
 ■ takes other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, others)  
 ■ takes more or for a longer time than directed ▶

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**  
 †This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark St. Joseph® Chewables  
**Manufactured by:**  
 Advance Pharmaceutical Inc.  
 Holtsville, NY 11742



NDC 17714-009-36

# CHILDREN'S ASPIRIN (NSAID)

1 1/4 gr (81 mg)

Pain Reliever

36 TABLETS

Chewable

Orange Flavor



**SEE CARTON FOR COMPLETE DRUG FACTS**

**Use** temporarily relieves minor aches and pains

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include ■ hives ■ asthma (wheezing) ■ facial swelling ■ shock

**Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood-thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, others) ■ takes more or for a longer time than directed

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ this product does not contain directions or complete warnings for adult use ■ administer only under adult supervision ■ if possible, use weight to dose, otherwise use age ■ repeat dose every 4 hours while symptoms last ■ do not give more than 5 doses in 24 hours ■ drink water with each dose

**77-83 lbs. (11-under 12 yrs):** 4-6 tablets, **66-76 lbs. (9-under 11 yrs):** 4-5 tablets, **46-65 lbs. (6-under 9 yrs):** 4 tablets, **36-45 lbs. (4-under 6 yrs):** 3 tablets, **32-35 lbs. (3-under 4 yrs):** 2 tablets, **under 32 lbs. (under 3 yrs):** ask a doctor

Store at room temperature 15°-30°C (59°-86°F)

631-981-4600

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



LA0710

Lot No.:  
Exp. Date:

**NDC: 17714-009-36 - 36 COUNT**

**ASPIRIN**

aspirin tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:17714-009
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ORANGE (UNII: 5EVU04N5QU)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	

**Product Characteristics**

<b>Color</b>	orange	<b>Score</b>	score with uneven pieces
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	AP;009
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17714-009-36	36 in 1 BOTTLE	04/15/1988	
1		1 in 1 CARTON; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	04/15/1988	

**Labeler** - Advance Pharmaceutical Inc. (078301063)**Registrant** - Advance Pharmaceutical Inc. (078301063)**Establishment**

Name	Address	ID/FEI	Business Operations
Advance Pharmaceutical Inc.		078301063	manufacture(17714-009)

Revised: 12/2024

Advance Pharmaceutical Inc.