

**HEB SPF 50 REFINE 50 SUNSCREEN WITH NIACINAMIDE AND COLLAGEN  
CONTINUOUS SUNSCREEN- avobenzone, homosalate, octisalate,  
octocrylene spray  
H.E.B**

-----

**HEB SPF 50 Refine 50 Sunscreen with Niacinamide and Collagen Continuous  
Sunscreen Spray**

***Active Ingredients***

Avobenzone 3.0%, Homosalate 10.0%, Octisalate 5.0%, Octocrylene 4.0%

***Purpose***

Sunscreen

***Uses***

- helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

***Warnings***

**For external use only**

**FLAMMABLE:**

- do not use near heat, flame or while smoking
- avoid long term storage above 104°F (40°C)

**Do not use**

- on damaged or broken skin.

**When using this product**

- keep out of eyes. Rinse with water to remove.
- do not puncture or incinerate.
- Contents under pressure.
- do not store at temperatures above 120°F.

**Stop use and ask a doctor if**

- rash occurs.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- apply liberally 15 minutes before sun exposure
- hold can 4-6 inches away from body, spray evenly to ensure complete coverage
- do not spray into face. Spray into hand and apply to face.
- use in well ventilated, but not windy areas
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying
  - at least every 2 hours
  - children under 6 months of age: Ask a doctor
  - **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
    - limit time in the sun, especially from 10 a.m. - 2 p.m.
    - wear long-sleeved shirts, pants, hats, and sunglasses

### **Other Information**

- protect the product in this container from excessive heat and direct sun
- may stain or damage some fabrics, materials or surfaces

### **☐ Inactive Ingredients**

SD ALCOHOL 40-B, BUTYLOCTYL SALICYLATE, FRAGRANCE, ACRYLATES/OCTYLACRYLAMIDE COPOLYMER, ALOE BARBADENSIS LEAF JUICE, BUTYLENE GLYCOL, SACCHAROMYCES/ PODOPHYLLUM PELTATUM FERMENT FILTRATE, DAUCUS CAROTA SATIVA (CARROT) ROOT EXTRACT, PANTHENOL, PERSEA GRATISSIMA (AVOCADO) OIL, VITIS VINIFERA (GRAPE) SEED OIL, SIMMONDSIA CHINENSIS (JOJOBA) SEED OIL, COCOS NUCIFERA (COCONUT) OIL, CAPRYLIC/CAPRIC TRIGLYCERIDE, GLYCERIN, STEAROXYTRIMETHYLSILANE, TOCOPHEROL, HYDROLYZED COLLAGEN, NIACINAMIDE.

### **Label**

Water Resistant (80 minutes)  
Paraben Free\*, PABA Free & Oxybenzone Free  
Contains Vitamins E & B5  
Hypoallergenic - Dermatologist Tested

Drug Facts	
<b>Active ingredients</b>	<b>Purpose</b>
Avobenzone 3% Homosalate 10% Octisalate 5% Octocrylene 4%	Sunscreen
<b>Uses</b> • helps prevent sunburn • if used as directed with other sun protection measures (see <i>Directions</i> ), decreases the risk of skin cancer and early skin aging caused by the sun	
<b>Warnings</b> For external use only	
<b>Flammable:</b> • do not use near heat, flame or while smoking • avoid long term storage above 104°F (40°C)	
Do not use • on damaged or broken skin.	
When using this product • keep out of eyes. Rinse with water to remove. • Keep away from face to avoid breathing it. • do not puncture or incinerate. Contents under pressure. • do not store at temperatures above 120° F.	
Stop use and ask a doctor if • rash occurs.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> • Spray liberally and spread evenly by hand 15 minutes before sun exposure • Hold can 4-6 inches away from the skin to apply • Do not spray directly into face. Spray into hands then apply to face • Do not apply in windy conditions • Use in a well-ventilated area • reapply: • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • children under 6 months of age: Ask a doctor • <b>Sun Protection Measures.</b> Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses	
<b>Other information</b> • protect the product in this container from excessive heat and direct sun • may stain or damage some fabrics, materials or surfaces.	
<b>Inactive ingredients</b> SD alcohol 40-8, butylacrylate/salicylate, acrylates/octylacrylamide copolymer, fragrance, hydrolyzed collagen, niacinamide, aloe barbadensis leaf juice, butylene glycol, saccharomyces/podophyllum peltatum ferment filtrate, oenothera carthagenensis (carrot) extract, water, panthenol, persea gratissima (avocado) oil, vitis vinifera (grape) seed oil, simmondsia chinensis (jojoba) seed oil, cocos nucifera (coconut) oil, caprylic/capric triglyceride, glycerin, stearoxydimethylsiloxane, xanthan.	
Questions or Comments? Call toll free 1-800-527-7731	

\*NO PARABENS SEPARATELY ADDED TO PRESERVE THIS PRODUCT

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TEXAS 78204



REFINE

50

Sunscreen with Niacinamide & Collagen

Niacinamide & Collagen help to improve skin appearance

BROAD SPECTRUM SPF 50



SUNSCREEN  
NET WT. 5.5 OZ (156 g)

## HEB SPF 50 REFINE 50 SUNSCREEN WITH NIACINAMIDE AND COLLAGEN CONTINUOUS SUNSCREEN

avobenzone, homosalate, octisalate, octocrylene spray

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-142
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	30 mg in 1 g
<b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	100 mg in 1 g
<b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	50 mg in 1 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	40 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>BOVINE TYPE I COLLAGEN</b> (UNII: FHJ3ATL51C)	
<b>PODOPHYLLUM</b> (UNII: 2S713A4VP3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>CARROT</b> (UNII: L56Z1JK48B)	
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>PANTHENOL</b> (UNII: WW9CM0O67Z)	
<b>JOJOBA OIL</b> (UNII: 724GKU717M)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ACRYLATE/ISOBUTYL METHACRYLATE/N-TERT-OCTYLACRYLAMIDE COPOLYMER (75000 MW)</b> (UNII: JU3XHR8VWK)	
<b>.ALPHA.-TOCOPHEROL, DL-</b> (UNII: 7QWA1RIO01)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>BUTYLOCTYL SALICYLATE</b> (UNII: 2EH13UN8D3)	
<b>GRAPE SEED OIL</b> (UNII: 930MLC8XGG)	
<b>COCONUT OIL</b> (UNII: Q9L0O73W7L)	
<b>AVOCADO OIL</b> (UNII: 6VNO72PFC1)	
<b>STEAROXYTRIMETHYLSILANE</b> (UNII: 9862TW94B2)	
<b>NIACINAMIDE</b> (UNII: 25X51I8RD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-142-37	156 g in 1 CAN; Type 0: Not a Combination Product	01/27/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	01/27/2022	

**Labeler** - H.E.B (007924756)