HEB FACE 50 CLEAR ZINC SUNSCREEN- octocrylene, zinc oxide lotion H.E.B.

HEB Face 50 Clear Zinc Sunscreen Lotion

Active ingredients

Octocrylene 4.0%, Zinc Oxide 5.0%

Purpose

Sunscreen

Uses

• helps prevent sunburn • if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use

on damaged or broken skin

When using this product

keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if

rash occurs

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply liberally 15 minutes before sun exposure
- reapply:
- after 80 minutes of swimming or sweating
- immediately after towel drying

- at least every 2 hours
- children under 6 months of age: Ask a doctor
- **Sun Protection Measures**. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- limit time in the sun, especially from 10 a.m. 2 p.m.
- wear long-sleeved shirts, pants, hats, and sunglasses

Other information

- protect the product in this container from excessive heat and direct sun
- may stain or damage some fabrics, materials or surfaces

Inactive ingredients

Water, Ethylhexyl Stearate, Diisopropyl Adipate, Cetyl Dimethicone, Cetyl PEG/PPG-10/1 Dimethicone, Hydrogenated Castor Oil, Polyglyceryl-4 Isostearate, Hexyl Laurate, Beeswax, Isohexadecane, Silica, Ethylhexylglycerin, Chlorphenesin, Fragrance, Triethoxycaprylylsilane, Dimethicone, Sodium Chloride, Aloe Barbadensis Leaf Juice, Sodium Ascorbyl Phosphate, Tocopherol.

Label





Clear Zinc **BROAD SPECTRUM** SPF 50 SUNSCREEN

UVA/UVB Protection Water Resistant (80 minutes) For Face, Nose and Ears





With Aloe & Vitamin E · Hypoallergenic

Drug Facts Active ingredients Purpose Octoorylene 4% Zinc Oxide 5% Sunscreen

- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use • on damaged or broken skin.

When using this product + keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if + rash occurs.

Keep out of reach of children. If swalltwed, get medical help or contact a Poison Control Center right away.

- Directions

 apply licerally to face, lips, ears and nose 15 minutes before sun exposure

 reapply:

 after 80 minutes of swimming or sweating

 immediately after towel drying

 all least every 2 hours

 seldeten uniter 6 months of ane; ask a doctor

- an east every 2 hours higher size ask a doctor
 Sun Protection Measures, Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad-spectrum SPF of 15 or higher and other sun protection measures including:
 I mit time in the sun, especially from 10 u.m. 2 u.m.
 weer long-sleeve shirts, pants, hats, and sunglasses

Other information

- protect this product from excessive heat and direct sun
 may stain or damage some fabrics, materials or surfaces

Inactive ingredients water, ethylnexyl stearate, disopropyl adipate, cetyl dimethicone, cetyl PEG/PPG-10/1 dimethicone, bydrogenated castor oil, polygłycecyl-4 isostearate, hepyl jeurate, beaswax, isohexadecane, silica, ethylnexylglycerin, chlarphenesin, fragrance, triethoxycapylystame, dimethicone, sodium chloride, aloe barbadensis leaf juice, sodium ascorbyl phosphate, tocopherol

Questions or Comments? Call toll free 1-800-527-7731

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TEXAS 78204







NET WT. 10Z (28.4 g)



(H-E-B)

Clear Zinc

Active Ingrithems: Occordyeria = %), cinc United 5 or 10 per sun protection measures (see Directions), decreases the risk of cancer and early skin aging caused by the sun Warnings: FOR EXTERNAL USE ONLY. Do not use • on damaged or broken skin, When using this product keep out of eyes. Rinse with water to remove. Stop use and ask a doctor if rash occurs. Keep out of reach of children, if swallowed, get medical help or contact a Poison Control Center right away. brokens kin, When using our product coop ducture year map and the product coop ducture year map and the product from excessive heat and direct sun. Directions: apply liberally to face, lips, ears and nose 15 minutes before sun exposure - reapply a ther 80 minutes of swimming or sweating - immediately after towed drying - at least every 2 hours: - children under 6 months of age, ask a doctor: - Sun Protection Measures, Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad-spectrum SPF of 15 or highter and other sun protection measures including * limit time in the sun, especially from 10 a.m. - 2 p.m. * wear long-sleeve shirts, pants, hats, and sunglesses

HEB FACE 50 CLEAR ZINC SUNSCREEN

octocrylene, zinc oxide lotion

Product Information

NDC:37808-151 **Product Type HUMAN OTC DRUG** Item Code (Source)

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	50 mg in 1 g	
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	40 mg in 1 g	

Inactive Ingredients	
Ingredient Name	Strength
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 5) (UNII: 035JKJ76MT)	
ISOHEXADECANE (UNII: 918X1OUF1E)	
ETHYLHEXYL STEARATE (UNII: EG3PA2K3K5)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
CETYL DIMETHICONE 45 (UNII: IK315POC44)	
DIISOPROPYL ADIPATE (UNII: P7E6YFV72X)	
WHITE WAX (UNII: 7G1J5DA97F)	
CHLORPHENESIN (UNII: 1670DAL4SZ)	
HEXAMETHYLDISILOXANE (UNII: D7M4659BPU)	
HYDROGENATED CASTOR OIL (UNII: ZF94AP8MEY)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
SODIUM ASCORBYL PHOSPHATE (UNII: 836SJG51DR)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
.ALPHATOCOPHEROL, DL- (UNII: 7QWA1RIO01)	
WATER (UNII: 059QF0KO0R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1	NDC:37808-151- 09	28.4 g in 1 JAR; Type 0: Not a Combination Product	01/07/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	01/07/2020	

Labeler - H.E.B. (007924756)

Revised: 10/2024 H.E.B.