

**TRANSCEND NPE FREE- iodine liquid
IBA**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

TRANSCEND NPE FREE

ACTIVE INGREDIENT

Iodine..... 1.00 %
(Provides A Full 1.0% Available Iodine As Used)

INERT INGREDIENTS

Emollients (Triple Ingredient System)..... 12.00 %
Other Inerts..... 87.00 %
(Includes Iodine Stabilizers, Buffering Agents and Surfactants)
TOTAL: 100.00 %

pH of 5.5 at time of manufacture
This product contains highly effective levels of
free iodine for fast kill of mastitis causing organisms

RECOMMENDED PROCEDURES FOR TEAT SANITATION

POST-MILKING (POST-DIPPING)

Immediately after milking, dip all teats full length with TRANSCEND NPE FREE Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

WARNING

DO NOT DILUTE THIS PRODUCT.
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
PROTECT THIS PRODUCT FROM FREEZING.
NOT FOR HUMAN USE
CAUTION: KEEP OUT OF REACH OF CHILDREN.

AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

Transcend NPE Free

- Unique "Drip-Less" Formulation
- Forms A Full Barrier With Plug
- Easily Removed Before Milking
- Compatible With Popular Pre-Dip Products



NDC # 29983-1072-6

LESS DRIP
LESS WASTE

NPE
FREE

TRANSCEND

NPE
FREE

SANITIZING 1% IODINE BARRIER TEAT DIP

A PREMIUM, "TRUE" BARRIER TEAT DIP

CONTAINS A **UNIQUE, 12%** TRIPLE INGREDIENT SKIN CONDITIONING SYSTEM

ACTIVE INGREDIENT:


Iodine.....	1.00%
<small>(Provides A Full 1.0% Available Iodine As Used)</small>	
INERT INGREDIENTS:	
Emollients (Triple Ingredient System).....	12.00%
Other Inerts.....	87.00%
<small>(Includes Iodine Stabilizers, Buffering Agents and Surfactants)</small>	
TOTAL:	100.00%

pH of 5.5 at time of manufacture
This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

RECOMMENDED PROCEDURES FOR TEAT SANITATION
POST-MILKING (POST-DIPPING)
Immediately after milking, dip all teats full length with **TRANSCEND NPE FREE Teat Dip**. Ensure good coverage of all sides of the teats and teat orifices.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.



An Aid In The Prevention
Of New Intramammary
Infections

WARNING

DO NOT DILUTE THIS PRODUCT.
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
PROTECT THIS PRODUCT FROM FREEZING.

NOT FOR HUMAN USE

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.**

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 15 GALLONS (56.775 LITERS) IBA STOCK # 260852 1223W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

TRANSCEND NPE FREE

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1072
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	1 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1072-4	19.70 kg in 1 PAIL		
2	NDC:29983-1072-6	59.11 kg in 1 DRUM		
3	NDC:29983-1072-8	216.75 kg in 1 DRUM		
4	NDC:29983-1072-9	1083.5 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/10/2018	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 12/2025

IBA