

ACETAMINOPHEN- acetaminophen tablet
RAONBIO CORP

Acetaminophen and Ibuprofen (NSAID) tablets 250mg/125mg
Dual Action
Pain Reliever

Active ingredients (in each caplet)

Acetaminophen 250 mg

Ibuprofen 125 mg (NSAID**)

Purposes

Pain Reliever

Uses

■ temporarily relieves minor aches and pains due to:
■ headache ■ toothache ■ backache ■ menstrual cramps ■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert: may cause severe skin reactions.

Symptoms may include:

- skin reddening ■ blisters ■ rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock
- skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Warnings

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

- chest pain ■ trouble breathing ■ weakness in one part or side of body
- slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed

adults and children 12 years and over ■ take 2 caplets every 8 hours while symptoms persist

children under 12 years ■ ask a doctor

- do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20 to 25°C (68 to 77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, crospovidone, ferric oxide red, ferric oxide yellow, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, sodium lauryl sulfate, stearic acid and titanium dioxide

Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

PDP

size : 1+3/4 X 1+3/4 X 3+3/8
 ref # : PP180604B
 material : .016 SBS

COATING FREE AREA

20000004711
 700000003524

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Best Choice

DUAL ACTION

Ibuprofen (NSAID) Tablets and Acetaminophen and Aspirin Tablets

250 mg/125 mg PAIN RELIEVER

72 CAPLETS: CAPSULE-SHAPED TABLETS

Actual Size

Compare to the active ingredients of Advil® Dual Action

CONTAINS 2 MEDICINES

Best Choice 100% Guaranteed

PROUDLY DISTRIBUTED BY:
 ASSOCIATED WHOLESALE DRUGGISTS, INC.
 KANSAS CITY, KANSAS 64108
 MADE IN INDIA

For product information please visit:
 bestchoicebrand.com/66723

www.bestchoicebrand.com

COATING FREE AREA

200000004711
 700000003524

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Do not use with other medicines containing these ingredients unless your doctor tells you to.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Drug Facts (continued)

Keep out of reach of children. If you have ever had an allergic reaction to acetaminophen or any other pain reliever, do not take this product.

Warnings:

- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.
- Do not take more than 6 capsules in 24 hours, or less as directed by a doctor.

Lot

Exp.

COATING FREE AREA

COATING
FREE AREA



Drug Facts (continued)

Acetaminophen can cause liver damage if you take too much. Ibuprofen can cause stomach bleeding, ulcers, and kidney problems if you take too much. Both can cause heart problems if you take too much. Do not take more than directed.

Directions

Take 2 caplets every 8 hours with meals and at bedtime. Do not take more than directed.

Other Information

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. If you are pregnant, get medical help right away.

Drug Facts (continued)

Acetaminophen can cause liver damage if you take too much. Ibuprofen can cause stomach bleeding, ulcers, and kidney problems if you take too much. Both can cause heart problems if you take too much. Do not take more than directed.

Directions

Take 2 caplets every 8 hours with meals and at bedtime. Do not take more than directed.

Other Information

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. If you are pregnant, get medical help right away.



COATING
FREE AREA

Drug Facts (continued)

Acetaminophen can cause liver damage if you take too much. Ibuprofen can cause stomach bleeding, ulcers, and kidney problems if you take too much. Both can cause heart problems if you take too much. Do not take more than directed.

Directions

Take 2 caplets every 8 hours with meals and at bedtime. Do not take more than directed.

Other Information

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. If you are pregnant, get medical help right away.

Questions or comments?
Call 1-877-776-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

PROUDLY DISTRIBUTED BY:
ASSOCIATED WHOLESALE DRUGGISTS, INC.
KANSAS CITY, KANSAS 64108

**Best Choice,
100% Guaranteed**

For product information please visit:
bestchoicebrand.com/66721

ref # : PP180604B
material : .016 SBS

Best Choice

DUAL ACTION

Ibuprofen (NSAID) Tablets

250 mg/125 mg

PAIN RELIEVER

18 CAPLETS

CAPULET-SHAPED TABLETS

Actual Size

Contains the active ingredients of Advil® Dual Action!

Drug Facts (continued)

Acetaminophen can cause liver damage if you take too much. Ibuprofen can cause stomach bleeding, ulcers, and kidney problems if you take too much. Both can cause heart problems if you take too much. Do not take more than directed.

Directions

Take 2 caplets every 8 hours with meals and at bedtime. Do not take more than directed.

Other Information

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. If you are pregnant, get medical help right away.

Drug Facts (continued)

Acetaminophen can cause liver damage if you take too much. Ibuprofen can cause stomach bleeding, ulcers, and kidney problems if you take too much. Both can cause heart problems if you take too much. Do not take more than directed.

Directions

Take 2 caplets every 8 hours with meals and at bedtime. Do not take more than directed.

Other Information

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. If you are pregnant, get medical help right away.

BEST CHOICE 66721 1023 GCH



Lot
Exp.

COATING
FREE AREA

size :1+3/4 X 1+3/4 X 3+3/8
 ref # :PP180604B
 material :.016 SBS



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70038-131
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	125 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE K30 (UNII: U725QWY32X)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
HYPROMELLOSE 2910 (15 MPA.S) (UNII: 36SFW2JZ0W)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	yellow ((Light yellow to yellow colored))	Score	no score
Shape	CAPSULE ((capsule shaped, biconvex))	Size	14mm
Flavor		Imprint Code	G;131
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70038-131-81	18 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2024	
2	NDC:70038-131-36	36 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2024	
3	NDC:70038-131-72	72 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA216592	01/01/2024	

Labeler - RAONBIO CORP (689517161)

