

INFANTS IBUPROFEN- ibuprofen suspension
Best Choice (Valu Merchandisers Company)

Ibuprofen Oral Suspension, USP

Drug Facts

Active ingredient (in each 1.25 mL)

Ibuprofen 50 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL = milliliter

- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every **6 to 8 hours**
- do not use more than **4 times a day**

| Dosing Chart | | |
|---------------------|------------------|------------------|
| Weight (lb) | Age (mos) | Dose (mL) |
| under 6 lbs | under 6 mos | ask a doctor |
| 12 to 17 lbs | 6 to 11 mos | 1.25 mL |
| 18 to 23 lbs | 12 to 23 mos | 1.875 mL |

Other information

- store between 20° to 25°C (68° to 77°F)
- **do not use if plastic bottle wrap imprinted “sealed for your protection” is broken or missing**

Inactive ingredients

carboxymethylcellulose sodium, citric acid anhydrous, glycerin, microcrystalline cellulose, natural and artificial berry fruit punch type flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, and xanthan gum.

Questions or Comments?

1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Principal Display Panel

Best Choice®

For Ages 6 Mos. to 23 Mos.

See New Warnings

COMPARE TO THE ACTIVE
INGREDIENT IN CONCENTRATED
MOTRIN® INFANTS' DROPS*

Infants'

Ibuprofen

Concentrated Ibuprofen

Oral Suspension, USP (NSAID)

50 mg per 1.25 mL

- Pain Reliever/Fever Reducer
- Lasts up to 8 hours
- Non-Staining
- Dye-Free
- Alcohol Free

BERRY FLAVOR

Use only with enclosed syringe

1 FL OZ (30 mL)

COATING FREE AREA

COATING FREE AREA

COATING FREE AREA

Best Choice

For Ages 6 Mos. to 23 Mos.

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL

Best Choice

For Ages 6 Mos. to 23 Mos.

See New Warnings

COMPARE TO THE ACTIVE INGREDIENT IN CONCENTRATED MOTRIN® INFANTS' DROPS*

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL



- Pain Reliever/Fever Reducer
- Lasts up to 8 hours
- Non-Staining
- Dye-Free
- Alcohol Free

BERRY FLAVOR

Use only with enclosed syringe

1 FL OZ (30 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 6 months to 23 months of age.

Drug Facts

Active ingredient (in each 1.25 mL)
Ibuprofen 50 mg (NSAID).....Pain reliever/fever reducer

Purpose
Pain reliever/fever reducer

Uses
temporarily:
relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include:
hives
facial swelling
asthma (wheezing)
shock
skin redness
rash
blister
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug
takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
takes more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

COATING FREE AREA

COATING FREE AREA

COATING FREE AREA

Best Choice

For Ages 6 Mos. to 23 Mos.

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL

Best Choice

For Ages 6 Mos. to 23 Mos.

See New Warnings

COMPARE TO THE ACTIVE INGREDIENT IN CONCENTRATED MOTRIN® INFANTS' DROPS*

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL



- Pain Reliever/Fever Reducer
- Lasts up to 8 hours
- Non-Staining
- Dye-Free
- Alcohol Free

BERRY FLAVOR

Use only with enclosed syringe

1 FL OZ (30 mL)

Drug Facts (continued)

Do not use
if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if
stomach bleeding warning applies to your child
child has a history of stomach problems, such as heartburn
child has problems or serious side effects from taking pain relievers or fever reducers
child has not been drinking fluids
child has lost a lot of fluid due to vomiting or diarrhea
child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
child has asthma
child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
under a doctor's care for any serious condition
taking any other drug

When using this product
take with food or milk if stomach upset occurs

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
child experiences any of the following signs of stomach bleeding:
feels faint
vomits blood
has bloody or black stools
has stomach pain that does not get better
child has symptoms of heart problems or stroke:
chest pain
trouble breathing
weakness in one part or side of body
slurred speech
leg swelling

Directions
this product does not contain directions or complete warnings for adult use
do not give more than directed
shake well before using
find right dose on chart below. If possible, use weight to dose; otherwise use age.
mL = milliliter
measure with the dosing device provided. Do not use with any other device.
dispense liquid slowly into the child's mouth, toward the inner cheek
if needed, repeat dose every 6 to 8 hours
do not use more than 4 times a day

COATING FREE AREA

COATING FREE AREA

COATING FREE AREA

Best Choice

For Ages 6 Mos. to 23 Mos.

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL

Best Choice

For Ages 6 Mos. to 23 Mos.

See New Warnings

COMPARE TO THE ACTIVE INGREDIENT IN CONCENTRATED MOTRIN® INFANTS' DROPS*

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL



- Pain Reliever/Fever Reducer
- Lasts up to 8 hours
- Non-Staining
- Dye-Free
- Alcohol Free

BERRY FLAVOR

Use only with enclosed syringe

1 FL OZ (30 mL)

Drug Facts (continued)

Do not use
if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if
stomach bleeding warning applies to your child
child has a history of stomach problems, such as heartburn
child has problems or serious side effects from taking pain relievers or fever reducers
child has not been drinking fluids
child has lost a lot of fluid due to vomiting or diarrhea
child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
child has asthma
child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
under a doctor's care for any serious condition
taking any other drug

When using this product
take with food or milk if stomach upset occurs

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
child experiences any of the following signs of stomach bleeding:
feels faint
vomits blood
has bloody or black stools
has stomach pain that does not get better
child has symptoms of heart problems or stroke:
chest pain
trouble breathing
weakness in one part or side of body
slurred speech
leg swelling

Directions
this product does not contain directions or complete warnings for adult use
do not give more than directed
shake well before using
find right dose on chart below. If possible, use weight to dose; otherwise use age.
mL = milliliter
measure with the dosing device provided. Do not use with any other device.
dispense liquid slowly into the child's mouth, toward the inner cheek
if needed, repeat dose every 6 to 8 hours
do not use more than 4 times a day

COATING FREE AREA

COATING FREE AREA

COATING FREE AREA

Best Choice

For Ages 6 Mos. to 23 Mos.

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL

Best Choice

For Ages 6 Mos. to 23 Mos.

See New Warnings

COMPARE TO THE ACTIVE INGREDIENT IN CONCENTRATED MOTRIN® INFANTS' DROPS*

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL



- Pain Reliever/Fever Reducer
- Lasts up to 8 hours
- Non-Staining
- Dye-Free
- Alcohol Free

BERRY FLAVOR

Use only with enclosed syringe

1 FL OZ (30 mL)

Drug Facts (continued)

Do not use
if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if
stomach bleeding warning applies to your child
child has a history of stomach problems, such as heartburn
child has problems or serious side effects from taking pain relievers or fever reducers
child has not been drinking fluids
child has lost a lot of fluid due to vomiting or diarrhea
child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
child has asthma
child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
under a doctor's care for any serious condition
taking any other drug

When using this product
take with food or milk if stomach upset occurs

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
child experiences any of the following signs of stomach bleeding:
feels faint
vomits blood
has bloody or black stools
has stomach pain that does not get better
child has symptoms of heart problems or stroke:
chest pain
trouble breathing
weakness in one part or side of body
slurred speech
leg swelling

Directions
this product does not contain directions or complete warnings for adult use
do not give more than directed
shake well before using
find right dose on chart below. If possible, use weight to dose; otherwise use age.
mL = milliliter
measure with the dosing device provided. Do not use with any other device.
dispense liquid slowly into the child's mouth, toward the inner cheek
if needed, repeat dose every 6 to 8 hours
do not use more than 4 times a day

COATING FREE AREA

COATING FREE AREA

COATING FREE AREA

Best Choice

For Ages 6 Mos. to 23 Mos.

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL

Best Choice

For Ages 6 Mos. to 23 Mos.

See New Warnings

COMPARE TO THE ACTIVE INGREDIENT IN CONCENTRATED MOTRIN® INFANTS' DROPS*

Infants' Ibuprofen

Concentrated Ibuprofen Oral Suspension, USP (NSAID)
50 mg per 1.25 mL



- Pain Reliever/Fever Reducer
- Lasts up to 8 hours
- Non-Staining
- Dye-Free
- Alcohol Free

BERRY FLAVOR

Use only with enclosed syringe

1 FL OZ (30 mL)

Drug Facts (continued)

Do not use
if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if
stomach bleeding warning applies to your child
child has a history of stomach problems, such as heartburn
child has problems or serious side effects from taking pain relievers or fever reducers
child has not been drinking fluids
child has lost a lot of fluid due to vomiting or diarrhea
child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
child has asthma
child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
under a doctor's care for any serious condition
taking any other drug

When using this product
take with food or milk if stomach upset occurs

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
child experiences any of the following signs of stomach bleeding:
feels faint
vomits blood
has bloody or black stools
has stomach pain that does not get better
child has symptoms of heart problems or stroke:
chest pain
trouble breathing
weakness in one part or side of body
slurred speech
leg swelling

Directions
this product does not contain directions or complete warnings for adult use
do not give more than directed
shake well before using
find right dose on chart below. If possible, use weight to dose; otherwise use age.
mL = milliliter
measure with the dosing device provided. Do not use with any other device.
dispense liquid slowly into the child's mouth, toward the inner cheek
if needed, repeat dose every 6 to 8 hours
do not use more than 4 times a day

INFANTS IBUPROFEN

ibuprofen suspension

| Product Information | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:63941-125 |
| Route of Administration | ORAL | | |

| Active Ingredient/Active Moiety | | |
|--|-------------------|------------------|
| Ingredient Name | Basis of Strength | Strength |
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 50 mg in 1.25 mL |

| Inactive Ingredients | |
|----------------------|----------|
| Ingredient Name | Strength |
| | |

| | |
|--|--|
| CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCROSE (UNII: C151H8M554) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Product Characteristics

| | | | |
|----------|----------------------------|--------------|--|
| Color | WHITE (white to off-white) | Score | |
| Shape | | Size | |
| Flavor | BERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:63941-125-24 | 1 in 1 CARTON | 10/02/2009 | |
| 1 | | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA079058 | 10/02/2009 | |

Labeler - Best Choice (Valu Merchandisers Company) (868703513)