

BERKLEY AND JENSEN OMEPRAZOLE- omeprazole tablet, delayed release BJWC

BJWC Omeprazole Delayed Release Tablets 20 mg Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting

- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-BJS-CLUB

Principal Display Panel

Compare to Prilosec OTC[®]

3 PACK

THREE 14-day courses of treatment

berkley jensen[®]

OMEPRAZOLE DELAYED RELEASE TABLETS, 20 mg

ACID REDUCER

ACTUAL SIZE

TREATS FREQUENT HEARTBURN!

24 HR

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



Compare to
Prilosec OTC®*

3 PACK

THREE 14-day
courses of treatment



OMEPRAZOLE
DELAYED RELEASE TABLETS, 20 mg
ACID REDUCER



ACTUAL
SIZE

**42
TABLETS**

THREE 14-DAY COURSES OF TREATMENT
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



TREATS FREQUENT
HEARTBURN!
24HR



9 3867 01801 9

Net Wt. 0.0728(2.06g)
BJ #42871

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole 20 mg.....Acid reducer

Use

- Treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight ■ Quit smoking

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

Inactive ingredients

canava wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-BJS-CLUB

Safety Feature -
Do not use if
printed tablet
blister unit is
open or torn.



Every Berkley Jensen
product comes with our
100% Money-Back Guarantee.

Made in Israel
Distributed by:
BJ's Wholesale Club, Inc.
350 Campus Drive
Marlborough, MA 01752

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

AA793 D3 C5

BERKLEY AND JENSEN OMEPRAZOLE

omeprazole tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-915
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
HYPROMELLOSE ACETATE SUCCINATE, UNSPECIFIED (UNII: A7ZHS2RJ34)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MONOETHANOLAMINE (UNII: 5KV86114PT)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	BROWN	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-915-55	3 in 1 CARTON	02/28/2008	
1		14 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA022032	02/28/2008	

Labeler - BJWC (159082692)

Revised: 2/2025

BJWC