

**NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled MEIJER, INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

**Pain reliever/Fever Reducer**

Cough Suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists more than 2 days, is accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

### **Other information**

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide\*, white ink

\*may contain this ingredient

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

Compare to Vicks® NyQuil® Cold & Flu LiquiCaps® active ingredients†

nitetime

Cold & Flu relief

Acetaminophen | 325 mg | Pain reliever / fever reducer

Dextromethorphan HBr | 15 mg | Cough suppressant

Doxylamine succinate | 6.25 mg | Antihistamine

MULTI-SYMPTOM RELIEF OF:

Aches, Fever, Sore throat, Cough, Runny Nose, Sneezing

ALCOHOL-FREE

Softgels

(\*\*Liquid-filled Capsules)

†This product is not manufactured or distributed by the Procter & Gamble Company. Vicks® , NyQuil® , and LiquiCaps® are registered trademark of the Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DIST. BY MEIJER

DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

[www.meijer.com](http://www.meijer.com)

**Package Label**

**meijer**

NDC 41250-743-24

Compare to Vicks® NyQuil®  
Cold & Flu LiquiCaps®  
active ingredients†

# nighttime cold & flu relief

**Acetaminophen | 325 mg | Pain reliever/Fever reducer**  
**Dextromethorphan HBr | 15 mg | Cough Suppressant**  
**Doxylamine succinate | 6.25 mg | Antihistamine**

**MULTI-SYMPATOM RELIEF OF:**

**Aches, Fever, Sore Throat, Cough, Runny Nose, Sneezing**

**ALCOHOL-FREE**

**24 Softgels\*\***

(\*\*) Liquid-filled Capsules



PER EVIDENCE: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

No.:  
Date:

**Drug Facts (continued)**

**Warnings**: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product ■ Allergy alert: Acetaminophen may cause severe skin reactions. ■ rash ■ skin reddening ■ blisters ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away. ■ **Sore throat warning**: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. ■ **Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ **Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus) ■ **Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin ■ **When using this product** ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ avoid alcoholic drinks ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

**Drug Facts (continued)**

**Directions** ■ do not take more than directed (see Overdose warning) ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ swallow whole; do not crush, chew, or dissolve ■ children under 12 years: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing ■ **Other information** ■ store between 15-30°C (59-86°F) ■ avoid excessive heat ■ titanium dioxide, white ink ■ polyethylene glycol, polyvinylpyrrolidone, purified water, sorbitan, sorbitol, FD&C yellow #10, FD&C blue #1, gelatin, glycerin ■ **Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Drug Facts**

**Active ingredients (in each softgel)**  
Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine

**Purposes**

■ temporarily relieves common cold and flu symptoms

**Uses**

■ fever ■ sore throat ■ headache ■ minor aches and pains ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ■ **If pregnant or breast-feeding**, ask a health professional before use. ■ **Keep out of reach of children.** ■ **Overdose warning**: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Liquid-filled Capsules

actual size



Lot #  
Exp.

**DIST. BY MEIJER  
DISTRIBUTION, INC.**  
GRAND RAPIDS, MI 49544  
www.meijer.com



\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

PID 467803



PLD-B41T FC005093

### MEIJER Nighttime Cold & Flu Relief

## NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-743
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P30;94A;AP017

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-743-24	24 in 1 CARTON	09/30/2018	12/27/2024
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/30/2018	12/27/2024

**Labeler** - MEIJER, INC. (006959555)

Revised: 9/2022

MEIJER, INC.