

**VICKS NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION- acetaminophen, diphenhydramine hcl, phenylephrine hcl powder, for solution  
Procter & Gamble Manufactura S de RL de CV**

-----  
**Vicks® NyQuil™ HOT REMEDY  
RAPID RELIEF  
MULTI-SYMPTOM  
COLD & FLU PLUS CONGESTION**

***Drug Facts***

**Active Ingredient (in each Packet)**

**Acetaminophen** 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Antihistamine, Cough Suppressant

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose
- sneezing

**Warnings**

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy Alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **Do not use more than directed**
- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

---

Age	Dose
Adults & children 12 yrs & over	one packet (dose) every 4 hours
Children under 12 yrs	Do not use

---

**Other information**

- **each packet contains:**potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- do not exceed 25°C.

## **Inactive ingredients**

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

## **Questions?**

**1-800-362-1683**

**Tamper Evident: Do not use if sachet is opened or torn**

**Keep outer package for complete product information.**

**MADE IN MEXICO**

**DIST. BY PROCTER & GAMBLE,**

**CINCINNATI OH 45202**

## **PRINCIPAL DISPLAY PANEL - 8 packet package**

MAX STRENGTH

Vicks® NyQuil™ HOT REMEDY

RAPID RELIEF MULTI-SYMPTOM

COLD & FLU + CONGESTION

**Acetaminophen**, Diphenhydramine HCl, Phenylephrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose, Cough

Nasal Congestion, Sinus Pressure

*Nighttime Relief*

HONEY LEMON FLAVORED

8 POWDER PACKETS

**MAX STRENGTH**

**VICKS**

# NyQuil™

## HOT REMEDY

### RAPID RELIEF MULTI-SYMPTOM

#### COLD & FLU + CONGESTION

Acetaminophen,  
Diphenhydramine HCl,  
Phenylephrine HCl

Headache, Fever,  
Sore Throat,  
Minor Aches  
& Pains

Sneezing,  
Runny Nose  
Cough

Nasal Congestion,  
Sinus Pressure

Nighttime Relief

**HONEY  
LEMON  
FLAVORED**

8 POWDER PACKETS

P&G  
www.pg.com  
vicks.com  
Patents: www.pg.com/patents

MADE IN MEXICO  
DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202  
90262782

**Questions?**  
1-800-382-1883

1570045751 6

**Drug Facts**

Active ingredients (in each Packet)	Purpose
Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose
- sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 6 doses in 24 hours, which is the maximum daily amount for this product.

• with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- Do not use more than directed
- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or occurs with rash or headache that lasts
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

**Age**

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

**Other information** each packet contains:

- potassium 10 mg
- phenylethanolamine: contains phenylalanine 68 mg per dose
- do not exceed 25°C

**Inactive ingredients:** croscellose potassium, aspartame, carboxymethylcellulose, citric acid, DSC Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

## VICKS NYQUIL HOT REMEDY RAPID RELIEF MULTI-SYMPTOM COLD AND FLU PLUS CONGESTION

acetaminophen, diphenhydramine hcl, phenylephrine hcl powder, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58933-551
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 g
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 5 g
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 5 g

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>TRIBASIC CALCIUM PHOSPHATE</b> (UNII: 91D9GV0Z28)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>CARBOXYMETHYLCELLULOSE</b> (UNII: 05JZI7B19X)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY (Lemon)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-551-01	5 g in 1 PACKET; Type 0: Not a Combination Product	03/22/2023	06/30/2027
2	NDC:58933-551-08	8 in 1 PACKAGE	03/22/2023	06/30/2027
2		5 g in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/22/2023	06/30/2027

**Labeler** - Procter & Gamble Manufactura S de RL de CV (812807550)