

GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN- acetaminophen tablet
GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN- acetaminophen tablet,
film coated
Unifirst First Aid Corporation

Green Guard® Maximum Strength Non-Aspirin

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not use more than directed (see overdose warning)

Adults and children (12 years and over)

- take 2 tablets every 6 hours while symptoms last.
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

Children under 12 years: Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch*, hypromellose*, polyethylene glycol*, povidone (K-30)*, pregelatinized starch*, purified water*, sodium starch glycolate, stearic acid, titanium dioxide*

*may contain**

Questions or comments? 1-800-869-6970

Maximum Strength Non-Aspirin relief

Acetaminophen 500 mg/Acetaminofeno 500 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Compare Active Ingredient To Extra Strength Tylenol®

Green Guard®

Pull Out

Tire Para Abrir

Tylenol® is a Registered Trademark of McNeil Consumer Products

50 Packets of 2 Tablets

100 Tablets

Order #2224

Distributed by Green Guard® St Louis, MO 63045

<p>Información del medicamento (continuación)</p> <p>Diagnóstico y consulte al médico si:</p> <ul style="list-style-type: none"> el dolor empeora o perdura más de 10 días la fiebre empeora o perdura más de 3 días aparecen nuevos síntomas aparece hinchazón o enrojecimiento <p>Estos pueden ser señales de una enfermedad grave.</p> <p>Si está usted embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.</p> <p>Mantenga este y todos los medicamentos fuera del alcance de los niños.</p> <p>Advertencia de sobredosis: En caso de ocurrir una sobredosis, comunique de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222). La información médica es crítica tanto para los adultos como para los niños aún cuando usted no sienta síntomas.</p> <p>Instrucciones</p> <p>■ no consuma más de lo que se indica (ver advertencia de sobredosis)</p> <p>Adultos y niños:</p> <ul style="list-style-type: none"> ■ Tome 2 tabletas cada 6 horas mientras los síntomas perduren (de 12 años y más) ■ no tomar más de 12 tabletas en 24 horas, a menos que lo indique el médico ■ no administrar por más de 10 días a menos que lo indique el médico <p>Niños menores de 12 años: consultar al médico</p> <p>Otra información</p> <ul style="list-style-type: none"> ■ almacenamiento: conservarse a temperatura ambiente 59-86°F (15-30°C) ■ impaguetado con sellado evidente ■ no use ningún paquete abierto o roto <p>Ingredientes inactivos</p> <p>amidon de maíz*, "hypromellose", polietilenglicol*, povidona (K-30)*, almidón pregelatinizado*, agua purificada*, almidón glicolato sódico, ácido estearico, dióxido de titanio*</p> <p>* puede contener</p> <p>¿Tiene alguna pregunta o comentario? Llame al 1-800-869-6970</p> <p style="text-align: right;">Rev. 11/823 Mfg. 1,24,14</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Acetaminophen 500 mg Pain reliever/fever reducer</p> <p>Uses</p> <p>Temporarily relieves minor aches and pains due to:</p> <ul style="list-style-type: none"> headache muscular aches backache the common cold premenstrual and menstrual cramps <p>Temporarily reduces fever.</p> <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4,000 mg in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Always alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are allergic to acetaminophen or any of the inactive ingredients in this product. <p>Ask a doctor before use if you have: liver disease</p> <p>Ask a doctor or pharmacist before use if:</p> <ul style="list-style-type: none"> you are taking the blood thinning drug warfarin <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> you get worse or lasts for more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p>
--	--

<p>Información del medicamento (continuación)</p> <p>Diagnóstico y consulte al médico si:</p> <ul style="list-style-type: none"> el dolor empeora o perdura más de 10 días la fiebre empeora o perdura más de 3 días aparecen nuevos síntomas aparece hinchazón o enrojecimiento <p>Estos pueden ser señales de una enfermedad grave.</p> <p>Si está usted embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.</p> <p>Mantenga este y todos los medicamentos fuera del alcance de los niños.</p> <p>Advertencia de sobredosis: En caso de ocurrir una sobredosis, comunique de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222). La información médica es crítica tanto para los adultos como para los niños aún cuando usted no sienta síntomas.</p> <p>Instrucciones</p> <p>■ no consuma más de lo que se indica (ver advertencia de sobredosis)</p> <p>Adultos y niños:</p> <ul style="list-style-type: none"> ■ Tome 2 tabletas cada 6 horas mientras los síntomas perduren (de 12 años y más) ■ no tomar más de 12 tabletas en 24 horas, a menos que lo indique el médico ■ no administrar por más de 10 días a menos que lo indique el médico <p>Niños menores de 12 años: consultar al médico</p> <p>Otra información</p> <ul style="list-style-type: none"> ■ almacenamiento: conservarse a temperatura ambiente 59-86°F (15-30°C) ■ impaguetado con sellado evidente ■ no use ningún paquete abierto o roto <p>Ingredientes inactivos</p> <p>amidon de maíz*, "hypromellose", polietilenglicol*, povidona (K-30)*, almidón pregelatinizado*, agua purificada*, almidón glicolato sódico, ácido estearico, dióxido de titanio*</p> <p>* puede contener</p> <p>¿Tiene alguna pregunta o comentario? Llame al 1-800-869-6970</p> <p style="text-align: right;">Rev. 11/823 Mfg. 1,24,14</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Acetaminophen 500 mg Pain reliever/fever reducer</p> <p>Uses</p> <p>Temporarily relieves minor aches and pains due to:</p> <ul style="list-style-type: none"> headache muscular aches backache the common cold premenstrual and menstrual cramps <p>Temporarily reduces fever.</p> <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4,000 mg in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Always alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are allergic to acetaminophen or any of the inactive ingredients in this product. <p>Ask a doctor before use if you have: liver disease</p> <p>Ask a doctor or pharmacist before use if:</p> <ul style="list-style-type: none"> you are taking the blood thinning drug warfarin <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> you get worse or lasts for more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p>
--	--

GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-170
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	44;148
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-170-33	50 in 1 BOX	02/01/2023	
1	NDC:47682-170-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-170-48	125 in 1 BOX	02/01/2023	
2	NDC:47682-170-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2023	

GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-619
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics

Color	white ((white to off white))	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	G552
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-619-33	50 in 1 CARTON	02/06/2024	
1	NDC:47682-619-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-619-48	125 in 1 CARTON	02/06/2024	
2	NDC:47682-619-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/06/2024	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(47682-170, 47682-619)

Establishment

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(47682-170, 47682-619)

Revised: 3/2024

Unifirst First Aid Corporation