

GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN- acetaminophen tablet
GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN- acetaminophen tablet,
film coated
Unifirst First Aid Corporation

Green Guard® Maximum Strength Non-Aspirin

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not use more than directed (see overdose warning)

Adults and children (12 years and over)

- take 2 tablets every 6 hours while symptoms last.
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

Children under 12 years: Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch*, hypromellose*, polyethylene glycol*, povidone (K-30)*, pregelatinized starch*, purified water*, sodium starch glycolate, stearic acid, titanium dioxide*

*may contain**

Questions or comments? 1-800-869-6970

Maximum Strength Non-Aspirin relief

Acetaminophen 500 mg/Acetaminofeno 500 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Compare Active Ingredient To Extra Strength Tylenol®

Green Guard®

Pull Out

Tire Para Abrir

Tylenol® is a Registered Trademark of McNeil Consumer Products

50 Packets of 2 Tablets

100 Tablets

Order #2224

Distributed by Green Guard® St Louis, MO 63045

Maximum Strength Non-Aspirin relief
Acetaminophen 500 mg/Acetaminofeno 500 mg
Pain Reliever/Fever Reducer
 Alivia el Dolor/Reduce la Fiebre

GREEN GUARD
 Distributed by Green Guard®, St Louis, MO 63045

Información del medicamento (continuación)
 Diga de inmediato a su médico si:
 ■ el dolor empeora o perdura más de 10 días
 ■ la fiebre empeora o perdura más de 3 días
 ■ aparecen nuevos síntomas
 ■ aparece hinchazón o enrojecimiento
 Estos pueden ser señales de una enfermedad grave.
 Si está usted embarazada o en periodo de lactancia, consulte la opinión de un médico antes de consumir este producto.
 Mantenga este y todos los medicamentos fuera del alcance de los niños.
Advertencia de sobredosis: En caso de ocurrir una sobredosis, comience de inmediato con un médico a hacer a un centro de control de envenenamiento (1-800-222-1222). La inmediata atención médica es crítica tanto para los adultos como para los niños aún cuando usted no note señales o síntomas.
Instrucciones
 ■ no consuma más de lo que se indica (ver advertencia de sobredosis)
Adulto y niño:
 ■ Tome 2 tabletas cada 6 horas mientras los síntomas perduren
 ■ no tomar más de 2 tabletas en 24 horas, a menos que lo indique el médico
 ■ no administrar por más de 10 días a menos que lo indique el médico
 Niños menores de 12 años: consultar al médico
Otra información
 ■ almacenamiento: conservarse a temperatura ambiente 59°-86°F (15°-30°C)
 ■ empacado con sellado evidente
 ■ no use ningún paquete abierto a otro
Ingredientes inactivos
 "hidróxido de metil", "hidroxido de sodio", "polietilenglicol", "propileno glicol", "almidón pregelatinizado", "agua purificada", "almidón glicolato sódico", "ácido estearico", "dióxido de titanio"
 *verde contener
 ¿Tiene alguna pregunta o comentario? Llame al 1-800-803-6970
 Rev. 11/02 Mfg. 1,24,44

Información del medicamento
Ingrediente activo (en cada tableta) Propiedad
 Acetaminofeno 500 mg Alivia el dolor/Reduce la fiebre
Usos
 ■ para el alivio temporal de leves dolores y malestares asociados con:
 ■ dolor de cabeza ■ dolores musculares
 ■ dolores de la artritis ■ dolor de espalda
 ■ dolor de muelas ■ dolor de dientes
 ■ dolores premenstruales y menstruales
 ■ reduce la fiebre de manera temporal
Advertencias
 Advertencia con respecto al hígado: Este producto contiene acetaminofeno. Dado que el hígado puede ocurrir si usted consume:
 ■ más de 4,000 mg en 24 horas, la cual es la máxima dosis recomendada
 ■ este producto con cualquier otro medicamento que contenga acetaminofeno
 ■ 3 o más bebidas alcohólicas diariamente mientras toma este producto
Información del medicamento (continuación)
 Alerta a las alergias: El acetaminofeno puede causar reacciones alérgicas en la piel. Las cuales pueden incluir: ■ enrojecimiento de la piel ■ ampollas ■ erupción
 Si se presenta una reacción alérgica en la piel, suspenda su consumo de inmediato y obtenga atención médica.
 No se administre
 ■ con ningún otro producto que contenga acetaminofeno (recetado o sin receta), si no está seguro o si su medicamento contiene acetaminofeno consulte con un médico o farmacéutico.
 ■ no use si siente náuseas o vómitos o si tiene un cuadro de sus ingredientes reactivos en este producto.
 Consulte a un médico antes de administrarle al usted si tiene una enfermedad del hígado.
 Consulte con un médico o farmacéutico antes de administrarle si el medicamento varía para el adelgazamiento de la sangre.
 (continúa en la solapa interior de la caja)
MAXIMUM STRENGTH Non-Aspirin relief
 COMPARTIMIENTO INGRESANTE DE ESTA SUSTANCIA (Tylenol)
 Pull to Open Tire Para Abrir
Acetaminophen 500 mg/Acetaminofeno 500 mg
Pain Reliever/Fever Reducer
 Alivia el Dolor/Reduce la Fiebre
GREEN GUARD
 Distributed by Green Guard® St Louis, MO 63045
 50 Packets of 2 Tablets
 100 Tablets
 Order #2224

Drug Facts
Active ingredient (in each tablet) Purpose
 Acetaminophen 500 mg Pain reliever/fever reducer
Uses
 Temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis
 ■ backache ■ the common cold ■ toothache
 ■ premenstrual and menstrual cramps
 Temporarily reduces fever
Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Always alert: Acetaminophen may obscure other reactions. Symptoms may include: ■ skin redness ■ hives ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Drug Facts (continued)
 Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product
 Ask a doctor before use if you have ■ liver disease
 Ask a doctor or pharmacist before use if
 ■ you are taking the blood thinning drug warfarin
 Stop use and ask a doctor if
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
Drug Facts (continued)
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 ■ do not use more than directed (see overdose warning)
Adults and Children (12 years and over)
 ■ take 2 tablets every 6 hours while symptoms last
 ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor
 ■ do not use for more than 10 days unless directed by a doctor
Children under 12 years
 Ask a doctor
Other information
 ■ store at room temperature 59°-86°F (15°-30°C)
 ■ tamper-evident sealed packets
 ■ do not use any opened or torn packets
Inactive ingredients
 con starch*, hydroxide*, polyethylene glycol*, propylene glycol*, pregelatinized starch*, purified water*, sodium starch glycolate, stearic acid, titanium dioxide*
 *may contain
Questions or comments?
 1-800-803-6970
 RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Maximum Strength Non-Aspirin relief

Acetaminophen 500 mg/Acetaminofeno 500 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Compare Active Ingredient To Extra Strength Tylenol®

Green Guard®

Pull Out

Tire Para Abrir

Tylenol® is a Registered Trademark of McNeil Consumer Products

50 Packets of 2 Tablets

100 Tablets

Order #2224

Distributed by Green Guard® St Louis, MO 63045

GREEN GUARD
Distributed by Green Guard, St. Louis, MO 63045

Acetaminophen 500 mg
Pain Reliever/Fever Reducer
Alivia el Dolor/Reduce la Fiebre

Non-Aspirin
MAXIMUM STRENGTH

100 Tablets

Rev. 11/621 Mfg. 1,24,14

Información del medicamento (continuación)
Mantenga este y todos los medicamentos fuera del alcance de los niños.
Advertencia de edad: En caso de ocurrir una sobredosis, comuníquese de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222). La prescripción de aspirina médica es crítica tanto para los niños como para los adultos cuando usted no sabe qué hacer o si debe tomar.

Indicaciones
● No tome más de lo que se indica (ver advertencia de sobredosis).
● Tome 2 tabletas cada 6 horas mientras los síntomas persisten (de 12 años y mayor).
● No tomar más de 6 tabletas en 24 horas, a menos que lo indique el médico.
● No administrar por más de 10 días a menos que lo indique el médico.

Otra información
● almacenamiento: conservarse a temperatura ambiente 59°-86°F (15°-30°C)
● empaquetado con sellado evidente
● no usar después de la fecha de caducidad o rotura

Ingredientes inactivos
almidón de maíz*, hipromelosa*, polietilenglicol*, povidona (K-30)*, almidón pregelatinizado*, agua purificada*, almidón glicolato sódico, ácido estearico, dióxido de titanio*

*Tiene alguna pregunta o comentario? Llame al 1-800-663-6370

Drug Facts
Active ingredient (in each tablet) Purpose
Acetaminophen (500 mg) Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
● headache ● muscle aches ● minor pain of arthritis ● backache ● the common cold ● toothache
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
● more than 4,000 mg in 24 hours, which is the maximum daily amount
● with other drugs containing acetaminophen
● 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
● skin redness ● hives ● rash ● if a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
Do not use:
● with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
● if you are allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if you have:
● liver disease
● you are taking the blood thinning drug warfarin
Stop use and ask a doctor if:
● you get worse or lasts for more than 3 days
● fever gets worse or lasts for more than 3 days
● new symptoms occur
● redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Drug Facts (continued)
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
● do not use more than directed (see overdose warning)
Adults and Children (12 years and over):
● take 2 tablets every 6 hours while symptoms last.
● do not take more than 6 tablets in 24 hours, unless directed by a doctor.
● do not use for more than 10 days unless directed by a doctor.
Ask a doctor.
Children under 12 years:
● ask a doctor.

Other information
● use at room temperature 59°-86°F (15°-30°C)
● larger evident unopened packets
● do not use any opened or torn packets

Inactive ingredients
corn starch*, hypromellose*, polyethylene glycol*, povidone (K-30)*, pregelatinized starch*, purified water*, sodium starch glycolate, stearic acid, titanium dioxide*

Questions or comments?
1-800-663-6370

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-170
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	44;148
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:47682-170-33	50 in 1 BOX	02/01/2023	
1	NDC:47682-170-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-170-48	125 in 1 BOX	02/01/2023	
2	NDC:47682-170-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2023	

GREEN GUARD MAXIMUM STRENGTH NON-ASPIRIN

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-619
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics

Color	white ((white to off white))	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	G552
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-619-33	50 in 1 CARTON	02/06/2024	
1	NDC:47682-619-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-619-48	125 in 1 CARTON	02/06/2024	
2	NDC:47682-619-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-619-99	2 in 1 PACKET; Type 0: Not a Combination Product	02/06/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/06/2024	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(47682-170, 47682-619)

Establishment

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(47682-170, 47682-619)

Revised: 9/2025

Unifirst First Aid Corporation