

MEDI-FIRST PLUS EXTRA STRENGTH NON ASPIRIN- acetaminophen tablet
MEDIQUE EXTRA STRENGTH APAP- acetaminophen tablet
MEDI-FIRST EXTRA STRENGTH NON ASPIRIN- acetaminophen tablet
Unifirst First Aid Corporation

Unifirst First Aid Acetaminophen 500 mg

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if

- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**

Adults and children: (12 years and over)

- take 2 tablets every 6 hours while symptoms last.
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

Children under 12 years: Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose*, polyethylene glycol*, povidone, pregelatinized starch*, sodium starch glycolate*, stearic acid*, titanium dioxide*

**may contain*

Questions or comments? 1-800-634-7680

Medique® Extra Strength APAP

Non-Aspirin Tablets

Tabletas sin Aspirina

Pain Reliever/Fever Reducer • Acetaminophen 500 mg

Marca Registrada de McNeil Consumer Products
 This Package is for Households without Young Children.
 Este Paquete Para Hogares Sin Niños Pequeños.
 Pull to Open
 Tire Para Abrir
 Tamper Evident Unit Dose Packets
 Empaquetado con Sellado Evidente en Dosis Unitarias
 250 Tablets
 (125 x 2)

MEDI-FIRST®
Extra Strength
Non-Aspirin

✓ Aches, Fever • Acetaminophen 500 mg
 Dolores, Fiebre • Acetaminofeno 500 mg

Pain Reliever/Fever Reducer
Alivia el dolor/Reduce la fiebre

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
 ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.

Tamper Evident Unit Dose Packets
 Empaquetado con Sellado Evidente en Dosis Unitarias

250 Tablets
(125 x 2)

MEDI-FIRST®
Extra Strength
Non-Aspirin

Pain Reliever/Fever Reducer
Alivia el dolor/Reduce la fiebre

Pull to Open
Tire the top

✓ Aches, Fever • Acetaminophen 500 mg
 Dolores, Fiebre • Acetaminofeno 500 mg

Compare active ingredient to:
Extra Strength Tylenol®
 Registered Trademark of McNeil Consumer Products
 Marca Registrada de McNeil Consumer Products

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
 ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.

Tamper Evident Unit Dose Packets
 Empaquetado con Sellado Evidente en Dosis Unitarias

804480X
Rev. 2/14/23
Fig. 124

Información del Medicamento

Ingrediente activo (por cada tableta)
 Acetaminofeno 500 mg **Alivia el dolor/Reduce la fiebre**

Propiedades

Uso

para el alivio temporal de leves dolores y molestias asociadas con:

- dolor de cabeza
- dolores leves de la artritis
- resaca común
- dolores premenstruales y menstruales
- dolor muscular
- dolor de espalda

reduzca la fiebre de manera temporal

Advertencias

Advertencia con respecto al hígado: Este producto contiene acetaminofeno. Dado que el hígado puede sufrir si usted consume:

- más de 4,000 mg en 24 horas, la cual es la máxima dosis recomendada
- este producto con cualquier otro medicamento que contenga acetaminofeno
- 3 o más bebidas alcohólicas diariamente mientras toma este producto

Alerta a las alergias: El acetaminofeno puede causar severas reacciones alérgicas en la piel. Las cuales pueden incluir:

- enrojecimiento de la piel
- ampollas
- erupción

Si se presenta una reacción alérgica en la piel, suspénda su consumo de inmediato y obtenga atención médica.

No se administre

- con ningún otro producto que contenga acetaminofeno (incluido si no está seguro si lo está seguro si su medicamento contiene acetaminofeno consulte con un médico o farmacéutico)
- si usted es alérgico al acetaminofeno o a cualquiera de los ingredientes listados en este producto

Consulte a un médico antes de administrarlo si usted tiene:

- enfermedad del hígado

Información del Medicamento (continuación)

Consulte con un médico o farmacéutico antes de administrarlo si:

- toma el medicamento warfarín para el adelgazamiento de la sangre

Deje de tomarlo y consulte al médico si

- el dolor empeora o persiste más de 10 días
- la fiebre empeora o persiste más de 3 días
- aparecen nuevos síntomas
- aparece inusual o empeoramiento

Estos pueden ser señales de una enfermedad grave.

Si está usted embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.

Mantenga este y todos los medicamentos fuera del alcance de los niños.

Advertencia de sobredosis: En caso de ocurrir una sobredosis, contacte de inmediato con un médico o llame a un centro de control de envenenamientos (1-800-222-1222). La inmediata atención médica es crítica tanto para los adultos como para los niños así como cuando usted no note señales o síntomas.

Instrucciones

- No consuma más de lo que se indica (ver advertencia de sobredosis)
- Tome 2 tabletas cada 6 horas mientras los síntomas persisten
- No tomar más de 6 tabletas en 24 horas, o menos que lo indique el médico
- No administrar por más de 10 días a menos que lo indique el médico

Niños menores de 12 años: consulte al médico

(continúa en el panel siguiente)

Información del Medicamento (continuación)

Otra información

- almacenamiento: consérvase a temperatura ambiente 69°-86°F (21°-30°C)
- empaquetado con sellado evidente
- no use ningún paquete abierto a otro.

Ingredientes inactivos

amplón de maíz, hypromelosa, polietilenglicol, polisorbo, amilón pregelatinizado, amilón glicosilado, ácido succínico, ácido succínico, ácido succínico, ácido succínico

¿Tiene alguna pregunta o comentario?
 Llame al 1-800-634-7680

Drug Facts

Active ingredient (in each tablet) **Purpose**
 Acetaminophen 500 mg Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- mild pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have: liver disease

Ask a doctor or pharmacist before use if:

- you are taking the blood thinning drug warfarin

Drug Facts (continued)

Stop use and ask a doctor if:

- you get worse or lasts more than 3 days
- you get worse or lasts more than 3 days
- you get worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)

Adults and children (12 years and over)

- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

Children under 12 years: Ask a doctor

Other information

- store at room temperature 69°-86°F (21°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, polyoxone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Other contents:


None

Questions or comments? 1-800-634-7680

Retain carton for complete product information

Medi-First® Plus Extra Strength Non-Aspirin
 Acetaminophen 500 mg
 Pain Reliever/Fever Reducer
 Pull to Open
 This Package is for Households without Young Children.
 Compare active ingredient to:
 Extra Strength Tylenol®
 Registered Trademark of McNeil Consumer Products
 Tamper Evident
 Unit Dose Packets
 100 Tablets

(50 x 2)



MEDI-FIRST PLUS EXTRA STRENGTH NON ASPIRIN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-174
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z 65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor		Imprint Code	44;148
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-174-33	50 in 1 BOX	02/01/2023	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-174-48	125 in 1 BOX	02/01/2023	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2023	

MEDIQUE EXTRA STRENGTH APAP

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-172
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor		Imprint Code	44;148
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-172-99	2 in 1 PACKET; Type 0: Not a Combination Product	02/01/2023	
2	NDC:47682-172-13	250 in 1 BOX	02/01/2023	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-172-33	50 in 1 BOX	02/01/2023	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-172-48	125 in 1 BOX	02/01/2023	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-172-64	12 in 1 BOX	02/01/2023	
5		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2023	

MEDI-FIRST EXTRA STRENGTH NON ASPIRIN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-173
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z 65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor		Imprint Code	44;148
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-173-99	2 in 1 PACKET; Type 0: Not a Combination Product	02/01/2023	
2	NDC:47682-173-13	250 in 1 BOX	02/01/2023	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-173-33	50 in 1 BOX	02/01/2023	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-173-48	125 in 1 BOX	02/01/2023	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-173-50	25 in 1 BOX	02/01/2023	
5		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2023	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment			
Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(47682-172, 47682-173, 47682-174)

Establishment			
Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(47682-172, 47682-173, 47682-174)