

FIBER LAXATIVE CALCIUM POLYCARBOPHIL- calcium polycarbophil tablet
Magno-Humphries, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Fiber Laxative Calcium Polycarbophil

MHL

NDC 54257-790-07

Compare to the active ingredient in Fibercon®*

FIBER LAXATIVE
CALCIUM POLYCARBOPHIL

- RELIEVES CONSTIPATION
- 280g of CALCIUM PER DOSE
- GENTLE ENOUGH TO USE EVERDAY

90 TABLETS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Active ingredient (in each tablet)

Calcium polycarbophil 625 mg
(Equivalent to 500 mg polycarbophil)

Purpose

Bulk-forming laxative

Uses

- relieves occasional constipation to help restore and maintain regularity
- this product generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- Do not use for more than 7 days unless directed by a doctor
- do not take more than 8 tablets in a 24 hour period unless directed by a doctor

Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

- take this product (child or adult dose) with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.

Adults and children over 12 years: 2 tablets once a day up to 4 times a day. **Children under 12 years:** ask a doctor

Other information

- **each tablet contains:** 140 mg calcium
- store at 20° to 25°C (68° to 77°F)
- do not refrigerate
- protect from humidity

Inactive ingredients

caramel, crospovidone, hypromellose, magnesium silicate, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silica, sodium lauryl sulfate, stearic acid.

Questions?

call toll-free **1-800-935-6737**

***This product is not manufactured or distributed by the owner of the registered trademark Fibercon®**

Distributed by: Magno-Humphries, Inc., OR 97223 U.S.A.

Visit our website: www.magno-humphries.com

*This product is not manufactured or distributed by the owner of the registered trademark Fibercon®

Distributed by: **Magno-Humphries, Inc., OR 97223 U.S.A.**

Visit our website: www.magno-humphries.com

Package Labeling:



Compare to the active ingredient in Fibercon®*

NDC 54257-790-07

FIBER LAXATIVE

CALCIUM POLYCARBOPHIL

- RELIEVES CONSTIPATION
- 280mg of CALCIUM PER DOSE
- GENTLE ENOUGH TO USE EVERYDAY



90 TABLETS





NDC 54257-790-07



Compare to the active ingredient in Fibercon®*

FIBER LAXATIVE
CALCIUM POLYCARBOPHIL

- RELIEVES CONSTIPATION
- 280mg of CALCIUM PER DOSE
- GENTLE ENOUGH TO USE EVERYDAY

90 TABLETS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each tablet)	Purpose
Calcium polycarbophil 625 mg.....	Bulk-forming laxative (Equivalent to 500 mg polycarbophil)

Uses • relieves occasional constipation to help restore and maintain regularity • this product generally produces bowel movement in 12 to 72 hours

Warnings Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have • abdominal pain, nausea, or vomiting • a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product • do not use for more than 7 days unless directed by a doctor • do not take more than 8 tablets in a 24 hour period unless directed by a doctor

Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

LIFT HERE

PEEL FOR DIRECTIONS

G7902-090-01-1

Drug Facts (continued)

Directions • take this product (child or adult dose) with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.

Adults and children over 12 years: 2 tablets once a day up to 4 times a day. **Children under 12 years:** ask a doctor

Other information • each tablet contains: 140 mg calcium • store at 20° to 25° C (68° to 77° F) • do not refrigerate • protect from humidity

Inactive ingredients caramel, crospovidone, hypromellose, magnesium silicate, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silica, sodium lauryl sulfate, stearic acid.

Questions? call toll-free 1-800-935-6737

*This product is not manufactured or distributed by the owner of the registered trademark Fibercon®.

Distributed by: **Magno-Humphries Labs** Tigard, OR 97223 U.S.A.

Visit our website: www.magno-humphries.com

FIBER LAXATIVE CALCIUM POLYCARBOPHIL

calcium polycarbophil tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54257-790	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W)		CALCIUM POLYCARBOPHIL	625 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CAMEL (UNII: T9D99G2B1R)				
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MAGNESIUM SILICATE (UNII: 9B9691B2N9)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
MINERAL OIL (UNII: T5L8T28FGP)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	white (off-white to cream)	Score	2 pieces	
Shape	OVAL	Size	19mm	
Flavor		Imprint Code	CPC339	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54257-790-07	90 in 1 BOTTLE; Type 0: Not a Combination Product	04/04/2017	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part334	04/04/2017		

Labeler - Magno-Humphries, Inc. (063251433)