

ASPIRIN 325- aspirin tablet, delayed release
Walgreen Company

Walgreens 44-691

Active ingredient (in each caplet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- shock
- hives
- facial swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 caplets every 4 hours not to exceed 12 caplets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

Walgreens

Compare to the active ingredient in
Bayer® Regular Strength Aspirin 325 mg
Safety Coated Caplets††

WALGREENS PHARMACIST RECOMMENDED†

NDC 0363-0691-12

Aspirin 325
325 mg / PAIN RELIEVER (NSAID)

Regular Strength

Enteric-Coated

ACTUAL SIZE

100
CAPLETS

HEART
HEALTH**

**Talk to your doctor or other healthcare provider
before using this product for your heart.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†Our pharmacists recommend the
Walgreens brand. We invite you to
compare to national brands.

††This product is not manufactured or
distributed by Bayer AG, owner of the
registered trademark Bayer® Regular
Strength Aspirin 325 mg Safety
Coated Caplets.

50844

ORG012269112

DISTRIBUTED BY: **WALGREEN CO.**
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens
100% SATISFACTION
GUARANTEED

walgreens.com
 ©2021 Walgreen Co.

No Print Area
Lot no. & Exp. Date



ACTUAL SIZE

HEART HEALTH

Compare to the active ingredient in Bayer® Regular Strength Aspirin 325 mg Safety Coated Caplets*

Aspirin 325
 325 mg / PAIN RELIEVER (NSAID) Regular Strength Enteric-Coated

100 CAPLETS

****Talk to your doctor or other healthcare provider before using this product for your heart.**

Walgreens

Compare to the active ingredient in Bayer® Regular Strength Aspirin 325 mg Safety Coated Caplets*



HEART HEALTH

NDC 0363-0691-12

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each caplet)
 Aspirin 325 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose
 Pain reliever

Uses
 For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings
 Reye's syndrome. Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ shock ■ asthma (wheezing)
 ■ hives ■ facial swelling ■ are age 60 or older

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older
 ■ take more or for a longer time than directed
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ have had stomach ulcers or bleeding problems
 ■ have 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 3 days
- fever gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

how2recycle.info



PAPER BOX
PLASTIC BOTTLE

ITEM 988239 W00000-0000-0
 REV0322

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



3 1917 119317 5

DISTRIBUTED BY: **WALGREEN CO.**
 200 WILMOT RD., DEERFIELD, IL 60015
walgreens.com
 ©2021 Walgreen Co.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not take more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 or 2 caplets every 4 hours not to exceed 12 caplets in 24 hours unless directed by a doctor
 ■ children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, polydextrose, polyethylene glycol, stearic wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-426-9391

50844 ORG012269112

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 3 days
- fever gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

DISTRIBUTED BY: **WALGREEN CO.**
 200 WILMOT RD., DEERFIELD, IL 60015
walgreens.com
 ©2021 Walgreen Co.

Walgreens 44-691

ASPIRIN 325

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0691
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	691
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0691-12	1 in 1 CARTON	07/01/2017	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a		

Combination Product			
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2017	

Labeler - Walgreen Company (008965063)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-0691)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0691)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0691)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(0363-0691)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-0691)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0691)