

FAMOTIDINE- famotidine tablet, film coated
Shopko Stores Operating Co., LLC

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Famotidine USP, 20 mg

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do Not Use

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

- store at 20° to 25°C (68° to 77°F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.
- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**

INACTIVE INGREDIENTS

Colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PATIENT INFORMATION

- **1 tablet** relieves heartburn due to acid indigestion
- Famotidine prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn - causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

PRINCIPAL DISPLAY PANEL

SHOPKO®

Compare to the Active Ingredient of Maximum Strength Pepcid® AC†

Maximum Strength

Acid Controller

Famotidine Tablets, USP 20 mg

Acid Reducer

Just One Tablet

Prevents & Relieves Heartburn Due to Acid Indigestion

25 Tablets

Manufactured by: Ohm Laboratories Inc.

5091040/0212

SHOPKO®

Maximum Strength

Acid Controller

Famotidine Tablets, USP 20 mg

Acid Reducer

SHOPKO®

Compare to the Active Ingredient of
Maximum Strength Pepcid® AC†

Maximum Strength

Acid Controller

Famotidine Tablets, USP 20 mg

Acid Reducer

Just One Tablet
Prevents & Relieves Heartburn Due to Acid Indigestion



25 Tablets

Tips for Managing Heartburn
■ Do not lie flat or bend over after eating ■ Do not wear tight-fitting clothing around the stomach
■ Do not eat before bedtime ■ Raise the head of your bed ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
■ Eat slowly and avoid big meals ■ If overweight, lose weight ■ Quit smoking

Expiration Date:

NON VARNISH

Batch No.

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.

See end panel for batch number and expiration date.

†This product is not manufactured or distributed by Johnson & Johnson • Merck Consumer Pharmaceuticals Co., distributor of Pepcid®, AC; Pepcid®, AC is a registered trademark of Merck & Co., Inc.




©2012 Shopko Stores
Operating Co., LLC
www.shopko.com
Manufactured by: Ohm Laboratories Inc.
1385 Livingston Avenue
North Brunswick, NJ 08902

0212

Drug Facts (continued)
Other information
■ store at 20° to 25° C (68° to 77° F)
■ protect from moisture
■ keep the carton. It contains important information.
Inactive ingredients colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide
Questions? call 1-800-406-7984

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
Do not use
■ with other acid reducers
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ if you have kidney disease, except under the advice and supervision of a doctor
Ask a doctor before use if you have
■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss
■ nausea or vomiting
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
Stop use and ask a doctor if
■ your heartburn continues or worsens ■ you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Directions
adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor



Drug Facts
Active ingredient (in each tablet) Famotidine USP, 20 mg
Purpose Acid reducer
Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking

FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:370 12-036
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	036
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:370 12-036-26	25 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA090283	07/23/2010	
------	------------	------------	--

Labeler - Shopko Stores Operating Co., LLC (023252638)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(37012-036)

Revised: 2/2013

Shopko Stores Operating Co., LLC