

PAIN RELIEVER- acetaminophen capsule, liquid filled Walgreens

Pain Reliever

Drug Facts

Active Ingredient

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours with other drugs
- containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

store between 20-25°C (68-77°F)

Inactive ingredients

FD & C Red # 40, gelatin, glycerin, polyethylene glycol 40, polyethylene glycol 400, povidone, purified water, sorbitan solution, titanium dioxide

Questions or comments?

1-800-373-6981

Compare to the active ingredient in Tylenol® Extra Strength

Pain Reliever

Acetaminophen 500 mg

Walgreens
Compare to the active ingredient in Tylenol® Extra Strength†

Mini Softgels **NEW**

Pain Reliever
ACETAMINOPHEN 500 mg / PAIN RELIEVER / FEVER REDUCER

Extra Strength

- Ultra concentrated mini softgels
- Dissolves quickly

50 MINI SOFTGELS

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER THE CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each softgel) Purpose
Acetaminophen Pain reliever/fever reducer
500 mg

Uses
temporarily relieves minor aches and pains due to:
 ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches
 ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning
This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours with other drugs ■ containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued under label)
 NDC 000-0000-00
 DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED walgreens.com ©2024 Walgreen Co.
 MADE IN INDIA ITEM 794816 W00000-0000-0
 W30RGMMYY-F
 3 11917 04362 3
PEEL HERE

Drug Facts (Continued)

- if you are allergic to acetaminophen or any of the inactive ingredients in this product
- Ask a doctor before use if you have liver disease
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> take 2 caplets every 6 hours while symptoms last do not take more than 6 caplets in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other Information
store between 20-25°C (68-77°F)

Inactive Ingredients
FD & C Red # 40, gelatin, glycerin, polyethylene glycol 40, polyethylene glycol 400, povidone, purified water, sorbitol solution, titanium dioxide

Questions or comments? 1-800-373-6981

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by JOHNSON & JOHNSON CONSUMER INC., owner of the registered trademark Tylenol®.

how2recycle.info

PLASTIC BOTTLE

PAIN RELIEVER

acetaminophen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1178
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SORBITAN (UNII: 6O92ICV9RU)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	SDP
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1178-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/26/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/26/2024	

Labeler - Walgreens (008965063)

Revised: 10/2025

Walgreens