SODIUM BICARBONATE- sodium bicarbonate tablet TWIN MED LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Sodium Bicarbonate 650 mg

Drug Facts

Active ingredient (in each tablet)

Sodium Bicarbonate 10 gr (650 mg)

Purpose

Antacid

Uses

relieves

- acid indigestion
- heartburn
- sour stomach
- upset stomach associated with these symptoms

Warnings

Do not use this product if you are on a sodium-restricted diet unless directed by a doctor. Do not take more than 24 tablets for adults up to 60 years of age (or 12 tablets for adults 60 years of age and older) in a 24 hour period nor use maximum dosage for more than 2 weeks, except under the advice and supervision of a physician. As with any drug, if you are pregnant or nursing a baby, seek advice of a health professional before using this product.

Stomach Warning

TO AVOID SERIOUS INJURY, DO NOT TAKE UNTIL TABLET IS COMPLETELY DISSOLVED IT IS VERY IMPORTANT NOT TO TAKE THIS PRODUCT WHEN OVERLY FULL FROM FOOD OR DRINK.

Consult a doctor if severe stomach pain occurs after taking this product.

Drug Interaction Precaution

Ask a physician or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. In case of

accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Directions

Adults

Take 1 tablet, dissolved in a glass of water, as needed. Maximum daily dose for adults up to 60 years of age is 24 tablets. Maximum daily dose for adults 60 years of age or older is 12 tablets. Dissolve completely in water before drinking.

DO NOT EXCEED RECOMMENDED DOSE. Not recommended for children.

Other information

- each tablet contains: sodium 178 mg (7.74 mEq)
- store at room temperature 15º-30ºC (59º-86°F) in a well-closed container as defined in the USP

Inactive ingredients

Microcrystalline cellulose and stearic acid, magnesium stearate

Questions or comments?

Questions ? call toll-free 1-844-912-4012

PRINCIPAL DISPLAY PANEL - 1000 Tablet Bottle Label

NDC 55681-331-10 Sodium Bicarbonate Tablets, USP

10 gr (650 mg)

ANTACID

Manufactured for: Twin Med LLC.	PR@CI PROC		G CHILDREN	Purpose Antacid	ed with these symptoms r adults 60 years of age spt under the advice and	acids may interact with	essistance or contact a	12 tablets in 24 hours 24 tablets in 24 hours		nesium stearate	
11333 Greenstone Ave Santa Fe Springs, CA 90670 ProCureProducts.com No Varnish Area	Sodium BICARBO	and the set of the set	FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN	ct)	 meson stormach = upset stormach associated with these symptoms addits up to 63 years of age (or 12 tablets for addits 60 years of age maximum dosage for more than 2 weeks, exect under the advise and 	a sodium restricted diet. Lee if yeu are taking a prescription drug, Arr	ore then 2 weeks. Ressional before use. Dential overdose, seek professional	than 2 weeks red in water prior to use ets every 4 hours, not more than is every 4 hours, not more than :	.74 mEq)	ine cellulose, stearic acid, mag	Questions or comments? Call to Tree 1-844-912-4012
NDC 55681-331-10 Item No. PCOTC103	Antacid	10 gr (650 mg) Item No. PCOTC103		gr (650 mg).	 m heartburn m sou 4 tabléts for adults up add nor use maximum 	H you have a sodium cist before use if yo	or if symptoms last more t willing, ask a health profess lithren. In case of accident rediately.	um dosage for more wed whole or dissol e and over: 1-2 tab s of age: 1-4 table	7 Sodium 178 mg (ure 15-30°C (59-86°F)	tits Microcrystal	nments? Cality
8 16516 01723 0	Relieves: • acid indigestion • heartburn • sour stomach • upset stomach	1000 TABLETS	THIS PACKAGE	Ctive Ingredient dium Bicarbonate 10 gr	1995 Nerves = acid indigestion Narmings o not take more than 24 nd oblech in a 24-hour pe	perveson or a prysocian sk a doctor before use sk a doctor or pharmai stain prescription drugs.	top use and ask a doct pregnant or breast-fee asp out of reach of chi lison Control Center Imr	Hrections do not use the maxim tablets may be swallo adults 60 years of ag adults under 60 year	Hther information each tablet contains: store at room temperat	nactive ingredie	Questions or comments?

	ate tablet								
Product Infor	mation								
Product Type		HUMAN OTC DF	۲UG	Item Code	e (Source)	NDC:55	681-331		
Route of Admini	stration	ORAL							
Active Ingredi	ent/Active	Moiety							
Ingredient Name						asis of trength	Strengt		
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - SODIU					S O DI UN BICARB	4	650 mg		
Inactive Ingre	dients								
		Ingredient	Name			S	Strength		
MICROCRYSTALLII	NE CELLULOS	SE (UNII: OP1R320	D61U)						
STEARIC ACID (UN									
MAGNESIIIM STEA	RATE (UNII: 7	0097M6I30)							
MAGNESION SILA									
	acteristics								
Product Chara		iite	Score			no score			
Product Chara Color	wł		Score Size			no score 10mm			
Product Chara Color Shape	wł	nite		Code					
Product Chara Color Shape Flavor	wł	nite	Size	Code		10mm			
Product Chara Color Shape Flavor Contains	wł	nite	Size	Code		10mm			
Product Chara Color Shape Flavor Contains Packaging # Item Code	wł RC	nite	Size Imprint (arketing Start Date	10mm S65	eting End Date		

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC monograph final	part331	10/10/2022					

Labeler - TWIN MED LLC (009579330)

Revised: 3/2023

TWIN MED LLC