

ARTHRITIS PAIN RELIEF- acetaminophen tablet, film coated, extended release

GERI-CARE PHARMACEUTICAL CORP

hst 265 (255)

Drug Facts

Active ingredient (in each caplet)

Acetaminophen USP, 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

Adults:

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

Under 18 years of age: ask a doctor

Other information

- store at 20 - 25° C (68 - 77° F). Avoid excessive heat 40° C (104° F).

Inactive ingredients

crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, sodium lauryl sulfate, stearic acid, titanium dioxide

Questions?

1-800-540-3765

NDC 57896-255-01

HEALTH STAR

ARTHRITIS PAIN RELIEF

ACETAMINOPHEN
EXTENDED-RELEASE TABLETS USP, 650 mg
PAIN RELIEVER/FEVER REDUCER
For the Temporary Relief of Minor Arthritis Pain
Lasts up to 8 hours. Use only as directed.

Compare to active ingredient in
8 Hour **TYLENOL**® Arthritis Pain**

**DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN**

100 Caplets* 650 mg each
(*Capsule-Shaped Tablets)

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS MISSING OR BROKEN.

Purpose
Acetaminophen USP, 650 mg. Pain reliever/fever reducer.

Drug Facts

Active ingredient (in each caplet)
Acetaminophen USP, 650 mg.

Uses
• temporarily relieves minor aches and pains due to:
• minor pain of arthritis • muscular aches • backache • toothache
• premenstrual and menstrual cramps • the common cold • headache
• temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
• more than 6 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions.
Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days
• new symptoms occur • redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not take more than directed (see overdose warning)
• take 2 caplets every 8 hours with water
• swallow whole; do not crush, chew, split or dissolve
• do not take more than 6 caplets in 24 hours
• do not use for more than 10 days unless directed by a doctor

Under 18 years of age: ask a doctor

Other information • store at 20 - 25°C (68 - 77°F).
Avoid excessive heat 40°C (104°F).

Inactive ingredients: crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, sodium lauryl sulfate, stearic acid, titanium dioxide

Questions or comments? 1-800-540-3765

** This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL**®.
REV 0228SP
DISTR. BY: **GERI-CARE**
PHARMACEUTICALS CORP.
1295 Towbin Ave
Lakewood, NJ 08701 3 57896 26501 4

ARTHRITIS PAIN RELIEF

acetaminophen tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-255
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
CROSPVIDONE (UNII: 2S7830E561)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	cor116
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-255-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076200	03/01/2023	12/31/2024

Labeler - GERI-CARE PHARMACEUTICAL CORP (611196254)**Registrant** - GERI-CARE PHARMACEUTICAL CORP (611196254)

Revised: 11/2024

GERI-CARE PHARMACEUTICAL CORP