

ASPIRIN - aspirin tablet
Velocity Pharma

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ASPIRIN 81mg

Active Ingredient
(in each tablet)

Aspirin 81 mg (NSAID)

Purpose

Pain Relief

Uses

- temporarily relieves minor aches and pains

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- Hives
- Asthma (wheezing)
- Shock
- Facial Swelling

Stomach Bleeding Warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- Are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- takes more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any pain reliever or fever reducer.

Ask a doctor before use if:

- The stomach bleeding warning applies to you
- You have history of stomach problems such as heartburn
- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You have asthma
- You are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for :

- Gout

- Diabetes
- Arthritis

Stop use and ask a doctor if

- you experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- New symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

These could be signs of serious conditions.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children 12 years and over:

For temporary relief of minor aches and pains, take 4 to 8 tablets every 4 hours while symptoms persist. Do not exceed 48 tablets in 24 hours unless directed by a doctor.

Children under 12 years of age: Do not use unless directed by a doctor.

Other Information

- Store at a controlled temperature between 62 and 77 degrees F

Inactive Ingredients

Anhydrous Lactose, Carnuba Wax Colloidal Silicon Dioxide, Crosscarmellose Sodium,FD&C Yellow #10 al lake, FD&C Yellow#6 al Lake, Iron Oxide Ochre, Methacrylic acid copolymer, Micro crystalline cellulose, Polysorbate 80, simethiocone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate.

Questions?

Call 1-855-314-1850

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Distributed by: Velocity Pharma LLC

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

TABLETS

60

Velocity

NDC 076168-009-10

Aspirin 81 mg (NSAID)

Low Dose

Safety Coated
Aspirin

Do not use if imprinted safety seal under cap is broken or missing.

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81mg (NSAID)	Pain Relief

Uses:
Temporarily relieves minor aches and pains.

Warnings: Keep outer carton for complete warnings and product information.

- Reye's Syndrome
- Allergy Alert
- Stomach bleeding warning
- Do not use
- Ask a doctor or pharmacist before use
- Stop use and ask a doctor
- Keep out of reach of children

Directions:
Adults and children 12 years and over:
For temporary relief of minor aches and pains, take 4 to 8 tablets every 4 hours while symptoms persist. Do not exceed 48 tablets in 24 hours unless directed by a doctor.
Children under 12 years of age:
Do not use unless directed by a doctor.

Questions? 1-855-314-1850

Distributed by
VELOCITY PHARMA LLC
226 B Sherwood Avenue
Farmingdale, NY 11735

VELASP81MG60L

Drug Facts (continued)

Ask a doctor before use if:

- The stomach bleeding warning applies to you
- You have history of stomach problems such as heartburn
- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You have asthma
- You are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- Gout
- Diabetes
- Arthritis

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- Feel faint
- Vomit blood
- Have bloody or black stools
- Have stomach pain that does not get better
- Allergic reaction occurs
- Ringing in the ears or loss of hearing occurs
- Pain gets worse or lasts more than 10 days
- New symptoms occur
- Redness or swelling is present. These could be signs of a serious condition.

If **pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during a delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Inactive Ingredients:

Anhydrous lactose, Carnuba Wax Colloidal Silicon Dioxide, Croscarmellose Sodium, FD&C Yellow # 10 al Lake, FD&C Yellow # 6 al Lake, Iron Oxide Ochre, Methacrylic Acid Copolymer, Micro Crystalline Cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate.

Directions

Adults and children 12 years and over:

For temporary relief of minor aches and pains, take 4 to 8 tablets every 4 hours while symptoms persist. Do not exceed 48 tablets in 24 hours unless directed by a doctor.

Children under 12 years of age:

Do not use unless directed by a doctor.

Other information: Store at a controlled temperature between 62 and 77 degrees F

Questions? 1-855-314-1850

60 TABLETS
81 mg each

60

Velocity

Aspirin 81 mg (NSAID)

Low Dose

Safety Coated Aspirin

Compare to the active ingredient in Bayer® Aspirin™

NDC 076168-009-10

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID) Pain Relief

Uses Temporarily relieves minor aches and pains.

Warnings

Reye's Syndrome. Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- Hives
- Facial Swelling
- Asthma (wheezing)
- Shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- Are age 60 or older
- Have had stomach ulcers or bleeding problems
- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any pain reliever or fever reducer. ▶

VELASP81MG60C

Enteric Coated
to help prevent
upset stomach

Distributed by

NDC: 76168-009-10 -60 COUNT

Enteric Coated to help prevent upset stomach

ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76 168-009
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
BROWN IRON OXIDE (UNII: 1N032N7MFO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	A
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76 168-009-10	60 in 1 BOTTLE		
1		1 in 1 CARTON		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/01/2011	

Labeler - Velocity Pharma (962198409)

Registrant - Velocity Pharma (962198409)

Revised: 10/2013

Velocity Pharma