

**ACID RELIEF- ranitidine tablet, film coated**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Acid Reducer Drug Facts**

**Active ingredient (in each tablet)**

Ranitidine 150 mg (as ranitidine hydrochloride 168 mg)

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water
- to **prevent** symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

**Other information**

- do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

**Inactive ingredients**

D&C yellow #10 aluminum lake, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, titanium dioxide, triacetin

**Questions or comments?**

**1-800-719-9260**

**Consumer Information****What you should know about MAXIMUM STRENGTH****Ranitidine Tablets, 150 mg****Acid Reducer**

(Please read all of this information before taking MAXIMUM STRENGTH Ranitidine Tablets, 150 mg. Save this leaflet for future reference.)


**What are MAXIMUM STRENGTH Ranitidine Tablets, 150 mg?**

- MAXIMUM STRENGTH Ranitidine Tablets, 150 mg contain 150 mg of ranitidine (as ranitidine hydrochloride, 168 mg), a medicine that doctors have prescribed more than 200 million times worldwide.

**Excellent Safety Record**

- The ingredient in MAXIMUM STRENGTH Ranitidine Tablets, 150 mg, ranitidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in MAXIMUM STRENGTH Ranitidine Tablets, 150 mg has been taken safely with

many frequently prescribed medications.



- MAXIMUM STRENGTH Ranitidine Tablets, 150 mg are sodium and sugar free.

### **What symptoms do MAXIMUM STRENGTH Ranitidine Tablets, 150 mg relieve and prevent?**

- MAXIMUM STRENGTH Ranitidine Tablets, 150 mg relieve and prevent heartburn associated with acid indigestion and sour stomach.
- Certain foods or beverages, and even lying down to sleep can cause heartburn associated with acid indigestion and sour stomach. It is normal for the stomach to produce acid, especially after consuming food or beverages. However, acid in the wrong place, such as the esophagus, or too much acid, can cause burning pain and discomfort.

### **How should I take MAXIMUM STRENGTH Ranitidine Tablets, 150 mg?**

- To **relieve** symptoms, swallow 1 tablet with a glass of water.
- To **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn.
- This medicine can be used up to twice daily (do not take more than 2 tablets in 24 hours).
- MAXIMUM STRENGTH Ranitidine Tablets, 150 mg should not be given to children under 12 years old unless directed by a doctor.
- **Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

### **How do MAXIMUM STRENGTH Ranitidine Tablets, 150 mg work?**

- MAXIMUM STRENGTH Ranitidine Tablets, 150 mg reduce the production of stomach acid. This is what makes MAXIMUM STRENGTH Ranitidine Tablets, 150 mg different from antacids, which neutralize the acid already in your stomach. Antacids do not reduce the production of acid.

### **Tips for managing heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables.
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking

- Raise the head of your bed
- Wear loose fitting clothing around your stomach

### **When should I see a doctor?**

- **Do not use**
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor
- **Ask a doctor before use if you have**
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- **Stop use and ask a doctor if**
- your heartburn continues or worsens
- you need to take this product for more than 14 days
- **If pregnant or breast-feeding**, ask a healthcare professional before use.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Questions or comments? 1-800-719-9260**

**BOTTLES: Bottle is sealed with printed foil under cap. Do not use if printed foil is open or torn.**

**BLISTERS: Do not use if the individual blister unit is open or torn.**

Distributed by

Perrigo®

Allegan, MI 49010

: 852B4 00 J2

### **Package/Label Principal Display Panel**

Compare to Zantac 150<sup>®</sup> active ingredient

MAXIMUM STRENGTH

acid relief

Ranitidine Tablets, 150 mg

Acid Reducer

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

65 Tablets (65 Doses) | actual size





**meijer**

NDC 41250-891-09

Compare to Zantac 150<sup>®</sup> active ingredient\*

**MAXIMUM STRENGTH**

# acid relief


Ranitidine Tablets, 150 mg  
Acid Reducer

**PREVENTS & RELIEVES HEARTBURN**  
associated with acid indigestion and sour stomach

**65 Tablets**  
(65 Doses)



actual size



LOT NO.

EXP.

052509 LE C3



P10 4442908

<b>Drug Facts</b>	<b>Purpose</b>
<b>Active ingredient (in each tablet)</b> Ranitidine 150 mg (as ranitidine hydrochloride 168 mg)	Acid reducer
<b>Uses</b> <ul style="list-style-type: none"> <li>relieve heartburn associated with acid indigestion and sour stomach</li> <li>prevent heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages</li> </ul>	
<b>Warnings</b> Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers. Do not use <ul style="list-style-type: none"> <li>if you have trouble or pain swallowing food, vomiting with blood, or blood or black stools. These may be signs of a serious condition. See your doctor.</li> <li>with other acid reducers</li> <li>if you have kidney disease, except under the advice and supervision of a doctor</li> </ul> Ask a doctor before use if you have <ul style="list-style-type: none"> <li>had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>heartburn with difficulty in swallowing, sweating or dizziness</li> <li>chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulder, or lightheadedness</li> <li>frequent chest pain</li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> </ul> Stop use and ask a doctor if <ul style="list-style-type: none"> <li>your heartburn continues or worsens</li> <li>you need to take this product for more than 14 days</li> </ul> If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
<b>Directions</b> <ul style="list-style-type: none"> <li>adults and children 12 years and over:               <ul style="list-style-type: none"> <li>to relieve symptoms, swallow 1 tablet with a glass of water</li> <li>to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn</li> </ul> </li> <li>can be used up to twice daily (do not take more than 2 tablets in 24 hours)</li> <li>children under 12 years: ask a doctor</li> </ul>	
<b>Other information</b> <ul style="list-style-type: none"> <li>do not use if printed on under bottle cap is open or torn</li> <li>store at 20°-25 °C (68°-77 °F)</li> <li>avoid excessive heat or humidity</li> <li>this product is sodium and sugar free</li> </ul>	

<b>Drug Facts (continued)</b>
<b>Inactive ingredients:</b> D&C yellow #10 aluminum lake, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, titanium dioxide, triacetin
<b>Questions or comments?</b> 1-800-719-0260

Read the directions, consumer information leaflet and warnings before use.

**MAXIMUM STRENGTH**  
**acid relief**  
Ranitidine Tablets, 150 mg  
Acid Reducer  
**PREVENTS & RELIEVES HEARTBURN**  
associated with acid indigestion and sour stomach

Keep the carton. It contains important information.  
 \*This product is not manufactured or distributed by Chatham, Inc.,  
 owner of the registered trademark Zantac 150\*

DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, MI 49544  
 www.meijer.com



## ACID RELIEF

ranitidine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-891
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

### Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>	BROWN ((Pinkish))	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10 mm
<b>Flavor</b>		<b>Imprint Code</b>	L852
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-891-09	1 in 1 CARTON	12/06/2018	
1		65 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-891-71	1 in 1 CARTON	01/08/2019	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-891-01	1 in 1 CARTON	03/27/2019	
3		95 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:41250-891-02	1 in 1 CARTON	05/28/2019	
4		24 in 1 BOTTLE; Type 0: Not a Combination Product		

5	NDC:41250-891-56	2 in 1 CARTON	05/28/2019	
5		1 in 1 CARTON		
5		95 in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA091429		12/06/2018	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 5/2019

Meijer Distribution Inc