

TUSSIN CF- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid
Discount Drug Mart

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 5 mL tsp)

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Phenylephrine HCL, USP 5 mg

Purpose

Cough Suppressant

Expectorant

Nasal Decongestant

Keep out of reach of children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses

- help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING.

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are
taking any other oral nasal decongestant or stimulant.

When using this product
do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding,
as a health professional before use.

Directions

- Do not take more than 6 doses in any 24 hour period
- This adult strength product is not intended for use in children under 12 years of age

Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use

Other information

- **Keep carton for full Directions for use.**
- store at 20°-25°C (68°-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients

Anhydrous citric acid, FD and red # 40, glycerin, menthol, natural and artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

DISCOUNT drug mart FOOD FAIR Tussin CF product label

*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH MULTI-SYMPTOM COLD

DISCOUNT

drug mart

FOOD FAIR

Tussin CF

Dextromethorphan HBr/ Guaifenesin / Phenylephrine HCL

NON-DROWSY

COUGH SUPPRESSANT

EXPECTORANT

NASAL DECONGESTANT

Relieves: Cough, Mucus, Nasal Congestion

Multi-Symptom Formula

For ages 12 and over

8 FL OZ (237 mL)

* This product is not manufactured or distributed bynPfizer, owner of the registered trademark Robitussin®

BX-040 LOT: EXP:

SATISFACTION GUARANTEED

IF DISSASIFIED , RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

DO NOT USE IF PRINTED SEAL UNDER CAP IS
TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)	Purpose
Dextromethorphan HBr, USP 10mg.....	Cough Suppressant
Guaifenesin, USP 100 mg.....	Expectorant
Phenylephrine HCl, USP 5 mg.....	Nasal Decongestant

Uses ■ help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
 ■ temporarily relieves these symptoms occurring with a cold:
 • nasal congestion
 • cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease
 ■ high blood pressure ■ thyroid disease ■ diabetes
 ■ trouble urinating due to an enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus)
 ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless
 ■ symptoms do not get better within 7 days or are accompanied by fever

■ cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ Do not take more than 6 doses in any 24 hour period
 ■ This adult strength product is not intended for use in children under 12 years of age

Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use

Other information ■ **Keep carton for full Direction for use.**
 ■ store at 20-25°C (68-77°F) ■ do not refrigerate
 ■ dosage cup provided ■ sodium 3 mg per teaspoonful

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®

BX-040



*COMPARE TO THE
ACTIVE INGREDIENTS IN
ROBITUSSIN® PEAK COLD
MULTI-SYMP TOM COLD



*COMPARE TO THE
ACTIVE INGREDIENTS IN
ROBITUSSIN® PEAK COLD
MULTI-SYMP TOM COLD

Tussin CF

Dextromethorphan HBr /
Guaifenesin /
Phenylephrine HCl
NON-DROWSY

**COUGH SUPPRESSANT
EXPECTORANT/
NASAL DECONGESTANT**

Relieves: Cough, Mucus,
Nasal Congestion

Multi-Symptom Formula
For ages 12 & over

8 FL OZ (237 mL)

Tussin CF

Dextromethorphan HBr /
Guaifenesin /
Phenylephrine HCl
NON-DROWSY

**COUGH SUPPRESSANT
EXPECTORANT/
NASAL DECONGESTANT**

Relieves: Cough, Mucus,
Nasal Congestion

Multi-Symptom Formula
For ages 12 & over

8 FL OZ (237 mL)

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE
PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR
DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG
MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



DISCOUNT
drug mart
FOOD FARE

*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD MULTI-SYMPOM COLD

Tussin CF

Dextromethorphan HBr /
Guaifenesin /
Phenylephrine HCl

**COUGH SUPPRESSANT
EXPECTORANT/
NASAL DECONGESTANT**

NON-DROWSY
Multi-Symptom Formula
For ages 12 & over
8 FL OZ (237 mL)

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)
Dextromethorphan HBr, USP 10 mg.....Cough Suppressant
Guaifenesin, USP 100 mg.....Expectorant
Phenylephrine HCl, USP 5 mg.....Nasal Decongestant

Uses ■ helps loosen phlegm (mucus) and bronchial secretions to drain bronchial tubes
■ temporarily relieves these symptoms occurring with a cold: ■ nasal congestion
■ cough due to minor throat and bronchial irritation.

Warnings
Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.
Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Stop use and ask a doctor ■ you get nervous, dizzy or sleepless
■ symptoms do not get better within 7 days or are accompanied by fever
■ cough lasts more than 7 days, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.
If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children.

Directions ■ do not take more than 6 doses in any 24 hour period

Age	Dose
Adults & children 12 years & over	2 teaspoons every 4 hours;
Children under 12 years	do not use

Other information ■ store at 20-25°C (68-77°F) ■ do not refrigerate
■ sodium 3 mg per teaspoonful ■ dosage cup provided ■ See carton for full labeling

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? 1-877-796-5944

* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®.

LOT:
EXP:

LR-070

TUSSIN CF

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-504
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K0OR)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-504-26	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/19/2014	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	11/19/2014		

Labeler - Discount Drug Mart (047741335)

Revised: 12/2018

Discount Drug Mart