SINUS CONGESTION AND PAIN SEVERE- acetaminophen, guaifenes in and phenylephrine hcl tablet, film coated

ARMY AND AIR FORCE EXCHANGE SERVICE

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Exchange Select 44-527C

Active ingredients (in each caplet)

Acetaminophen 325 mg Guaifenesin 200 mg Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer Expectorant Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a

doctor or pharmacist before taking this product.

• if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- heart disease
- thyroid disease
- diabetes
- liver disease
- high blood pressure
- difficulty in urination due to an enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse of lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs of symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15° 30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicone dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

exchange[|selectTM

Compare To
The Active Ingredients of
Tylenol® SINUS SEVERE*

Pseudoephedrine Free

SINUS

SEVERE CONGESTION & PAIN

Acetaminophen, Guaifenesin, Phenylephrine HCl Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

- Sinus Headache
- Nasal Congestion
- Sinus Pressure
- Mucus + Chest Congestion

DAY

NON-DROWSY

Actual Size

24 Caplets

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® SINUS SEVERE.

50844 REV0917E52708

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"

Manufactured For Your Military Exchanges Distributed by: LNK International, Inc. Hauppauge, NY 11788 1-800-426-9391



Drug Facts (continued)

DUCTINFO

Drug Facts

ot no. & Exp. Date

Corporation, owner of the registered trademark Tylenol® SINUS SEVERE.

50844 REV0917E52708

Compare To The Active Ingredients of Tylenol® SINUS SEVERE* Pseudoephedrine Free

exchange select

SEVERE CONGESTION & PAIN

Acetaminophen, Guaifenesin, Phenylephrine HCI Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant

- Sinus Headache
- Nasal Congestion
- Sinus Pressure
- Mucus + Chest Congestion

quality value

24 Caplets

 see end flap for expiration date and lot number 12,-30,C (26,-89,E)

■ store at 25°C (77°F); excursions permitted between BLISTER IS TORN OR BROKEN DO NOT USE IF OUTER PACKAGE IS OPENED OR ■ TAMPER EVIDENT: Other information

Drug Facts (continued)

acid, sucralose, talc, titanium dioxide stricon dioxide, sodium starch glycolate, stearic polyethylene glycol, polywnyl alcohol, povidone, maltodextrin, microcnystalline cellulose, crospovidone, flavor, magnesium stearate, Inactive ingredients comstach,

Actual Size

Orug Facts (continued)

- smoking, asthma, chronic bronchitis, or
- persistent or chronic cough such as occurs with condp tyst occurs with too much phiegm (mucus) Ask a doctor before use if you have

product or any of its ingredients

- It you have ever had an all ergic reaction to this or pharmacist before taking this product. prescription drug contains an MAOI, ask a doctor the MAOI drug. If you do not know if your Parkinson's disease), or for 2 weeks after stopping depression, psychiatric or emotional conditions, or
- oridase inhibitor (NAM) (certain drugs for u you are now taking a prescription monoanine qoctor or pharmadist. sure whether a drug contains acetaminophen, ask a
- (bisscription or nonprescription). If you are not мир зий один дила соитяший эсетяшиорини asu ton of

help right away.

ae

It a skin reaction occurs, stop use and seek medical ■ blisters ■ rash ■ skin reddening

reactions. Symptoms may include: Allergy alert: Acetaminophen may cause severe skin this product

- dhildren under 12 years: ask a doctor
- do not take more than 10 caplets in 24 hours swallow whole - do not crush, chew, or dissolve
 - fake 2 caplets every 4 hours
 - adults and children 12 years and over do not take more than directed

Directions

swodw/s children even if you do not notos any signs or medical attention is critical for adults as well as for Control Center (1-800-222-1222) right away. Prompt overdose, get medical help or contact a Poison Keep out of reach of children. In case of accidental professional before use

If pregnant or breast-feeding, ask a health

serious condition.

- headache that lasts. These could be signs of a
 - condu course pack or occurs with rash or uew symptoms occur
 - Ledness or swelling is present ■ fever gets worse or lasts more than 3 days
- lasts more than 7 days pain, nasal congestion, or cough gets worse or ■ uervousness, dizziness, or steepiessness occur


iue Area

B-0066E-527C-08-SS REV0917E52708

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts (continued) Questions or comments?

1-800-426-9391



Exchange Select 44-527C

SINUS CONGESTION AND PAIN SEVERE

acetaminophen, guaifenesin and phenylephrine hcl tablet, film coated

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-527	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg		
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg		
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		

Inactive Ingredients			
Ingredient Name	Strength		
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MALTO DEXTRIN (UNII: 7CVR7L4A2D)			
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)			
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)			
STARCH, CORN (UNII: O8232NY3SJ)			
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			

Product Characteristics				
Color	WHITE	Score	no score	

Shape	OVAL	Size	19 mm
Flavor	MINT	Imprint Code	44;527
Contains			

	Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:55301-527-08	2 in 1 CARTON	08/06/2005		
ı	1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part341	08/06/2005		

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(55301-527)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(55301-527)

Revised: 6/2019 ARMY AND AIR FORCE EXCHANGE SERVICE