

## **PRO NUMB- tattoo numbing spray spray**

### **Pro Numb Tattoo Numbing Spray LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

## **Pro Numb Tattoo Numbing Spray 120 mL**

### **Active Ingredient(s)**

Lidocaine HCl 5% Epinephrine HCl .01%

### **Purpose**

Topical Anesthetic

### **Use**

Temporarily takes away pain, itching, and swelling associated with anorectal disorders or pain-sensitive procedures.

### **Warnings**

External Use Only. Keep out of reach of children.

Do not use if you have • seizures or liver disease • if pregnant or breastfeeding • a known allergy or sensitivity occurs, discontinue use, and seek medical attention. If the condition intensifies or symptoms persist for more than seven days or clear up and occur again within a few days, Cease the use of this product and consult a doctor. Do not use in large quantities, Particularly over raw surfaces or blistered areas. Avoid contact with the eyes. If contact should occur, irrigate with large amounts of sterile saline and seek medical attention. Do not swallow, if swallowed, seek medical attention immediately.

When using this product, you may notice temporary whiting or redness of the skin where pro numb is applied.

Keep out of reach of children.

### **Directions**

A sensitivity test is encouraged before using. Spray a small amount on the broken area of the skin. Wait 60 seconds for the anesthetic to start working. Wash Pro Numb off completely before tattooing.

### **Other information**


Store in a cool dark place or refrigerate and keep out of reach of children. Discard the item after the expiration date is due.

### Inactive ingredients

purified RO water, Paraben-DU, Disodium EDTA, Citric acid, Optiphen.

Call: (800) 865-4965

### Package Label - Principal Display Panel



**Topical Anesthetic**  
**5% Lidocaine**

4FL OZ (120 ML)

**Drug Fact- For Use by Licensed Tattoo artists only**

<b>Active Ingredients</b>		<b>Purpose</b>
Lidocaine HCl	5%	Topical Anesthetic
Epinephrine HCl	.01%	Vasoconstriction

**Uses: External Use Only.**  
Temporarily takes away pain, itching, and swelling associated with anorectal disorders or pain-sensitive procedures.

**Warnings-Keep out of reach of children**  
**Do not use if you have**

- seizures or liver disease
- if pregnant or breastfeeding
- a known allergy or sensitivity occurs, discontinue use, and seek medical attention. If the condition intensifies or symptoms persist for more than seven days or clear up and occur again within a few days, Cease the use of this product and consult a doctor. Do not use in large quantities, Particularly over raw surfaces or blistered areas.

Avoid contact with the eyes. if contact should occur, irrigate with large amounts of sterile saline and seek medical attention.  
**DO NOT SWALLOW.** if swallowed, seek medical attention immediately.

**When using this product** you may notice temporary whitening or redness of the skin where pro numb is applied.

**Directions: Sensitivity test encouraged before use.**  
spray a small amount on the broken area of the skin. Wait 60 seconds for the anesthetic to start working. **Wash Pro Numb off completely before tattooing.**

**Inactive Ingredients:** purified RO water, Paraben-DU, Disodium EDTA, Citric acid, Optiphen.

**Other information:** Store in a cool dark place or refrigerate and keep out of reach of children. Discard after the expiration date.

**Questions?** Contact pro numb Distribution

**Distributed by:**  
Pro Numb Tattoo numbing spray LLC  
Melbourne, FL, 32907 (800) 865-4965  
[www.tattoonumbingsprays.com](http://www.tattoonumbingsprays.com)

120mL Spray NDC: 83389-111-11

## PRO NUMB

tattoo numbing spray spray

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83389-111
<b>Route of Administration</b>	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
EPINEPHRINE HYDROCHLORIDE (UNII: WBB047OO38) (EPINEPHRINE - UNII:YKH834O4BH)			EPINEPHRINE HYDROCHLORIDE	0.1 mg in 100 mL
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)			LIDOCAINE HYDROCHLORIDE ANHYDROUS	5 mg in 100 mL
Inactive Ingredients				
Ingredient Name				Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)				
WATER (UNII: 059QF0KO0R)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83389-111-11	120 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/12/2023	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other			04/12/2023	

**Labeler** - Pro Numb Tattoo Numbing Spray LLC (118964052)

Revised: 12/2025

Pro Numb Tattoo Numbing Spray LLC