# CHEWABLE LOW DOSE ASPIRIN- aspirin tablet, chewable TARGET CORPORATION

-----

### 334R-Target-11673-493-Chewable Low Dose Aspirin 81 mg

# **Drug Facts**

## Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

## **Purpose**

Pain reliever

#### Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for 81 mg Orange chewable aspirin

# Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

## Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

## Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

# If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **Directions**

- · drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to

exceed 48 tablets in 24 hours unless directed by a doctor

children under 12 years: consult a doctor

#### Other information


- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

### Inactive ingredients

butylated hydroxytoluene, dextrates, FD&C yellow #6 aluminum lake, flavor, microcrystalline cellulose, modified corn starch, pregelatinized starch, sodium saccharin, stearic acid

### Questions or comments?

### Call 1-800-910-6874





■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

■ drink a full glass of water with each dose

Directions

unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

definitely directed to do so by a doctor because it may cause problems in the If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless.

■ ringing in the ears or a loss of hearing occurs

**Drug Facts** (continued)

uew symptoms occur

■ redness or swelling is present ■ fever gets worse or lasts more than 3 days

■ you experience any of the following signs of stomach bleeding:

■ an allergic reaction occurs. Seek medical help right away. Stop use and ask a doctor if

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

■ you nave astnma

■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

■ you have a history of stomach problems, such as heartburn stomach bleeding warning applies to you

Ask a doctor before use if

■ if you have ever had an allergic reaction to this product or any of its ingredients ру а аостог

■ this product for at least 7 days after tonsillectomy or oral surgery unless directed

if you are allergic to aspirin or any other pain reliever/fever reducer

■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product

ipuproten, naproxen, or otners) ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin,

■ take a blood thinning (anticoagulant) or steroid drug

■ have had stomach ulcers or bleeding problems

changes in behavior with nauses and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert. Aspirin may cause a severe allergic reaction which may include:

Stomach bleeding warming: This product contains an MSAID, which may cause severe stomach bleeding warming: This product contains an MSAID, which may cause severe stomach bleeding. The chance is higher if you servere atomach bleeding. The chance is higher if you

pox or flu-like symptoms should not use this product. When using this product, if Reye's syndrome: Children and teenagers who have or are recovering from chicken

■ sak your doctor about other uses for 81 mg Orange chewable aspirin

■ for the temporary relief of minor aches and pains or as recommended by your doctor Sasn

Aspirin 18 mg (ASM) \*.....\* \*nonsteroidal anti-inflammatory drug

Active ingredient (in each chewable tablet)

Drug Facts

Other information

Call 1-800-910-6874

Questions or comments?

starch, sodium saccharin, stearic acid modified corn starch, flavor, pregelatinized

Inactive ingredients butylated

■ use by expiration date on package

hydroxytoluene, dextrates, FD&C yellow #6 aluminum lake, microcrystalline cellulose,

■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

'This product is not manufactured or distributed by Bayer AC, owner of the registered trademark Bayer® Chewable Aspirin.

Drug Facts (continued)

up&up

Purpose

81 mg Pain Reliever (NSAID)

Compare to active ingredient in Bayer® Chewable Aspirin<sup>†</sup>

094 01 0280 R00 C-002262-01-034-0000 Distributed by Target Corporation Minneapolis, MN 55403 TM & ©2024 Target Brands, Inc.


Love it or your money back

Satisfaction guaranteed

0 49319

Lot No.:

**Soating/Print Omit Area** 



# **CHEWABLE LOW DOSE ASPIRIN**

aspirin tablet, chewable

### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:11673-493

Route of Administration ORAL

### **Active Ingredient/Active Moiety**

Ingredient Name	<b>Basis of Strength</b>	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	AS PIRIN	81 mg

Inactive Ingredients			
Ingredient Name	Strength		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
MODIFIED CORN STARCH (1-OCTENYL SUCCINIC ANHYDRIDE) (UNII: 461P5CJN6T)			
STARCH, CORN (UNII: O8232NY3SJ)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

DEXTRATES (UNII: G263MI44RU)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)

Product Characteristics			
Color	orange	Score	no score
Shape	ROUND (bi-convex)	Size	8mm
Flavor	ORANGE	Imprint Code	TCL334
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-493- 19	3 in 1 CARTON	01/15/2024	
1	NDC:11673-493- 63	36 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/15/2024	

# **Labeler - TARGET CORPORATION (006961700)**

# **Registrant - TIME CAP LABORATORIES, INC. (037052099)**

Establishment					
Name	Address	ID/FEI	<b>Business Operations</b>		
TIME CAP LABORATORIES, INC		037052099	manufacture(11673-493)		

Revised: 1/2025 TARGET CORPORATION