ANTI DIARRHEAL- loperamide hcl tablet ARMY AND AIR FORCE EXCHANGE SERVICE

Exchange Select 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

drink plenty of clear fluids to help prevent dehydration caused by diarrhea

• find right dose on chart. If possible, use weight to dose; otherwise, use age.

| adults and children 12 years and | 2 caplets after the first loose stool; 1 caplet after each |
|----------------------------------|---|
| over | subsequent |
| over | loose stool; but no more than 4 caplets in 24 hours |
| | 1 caplet after the first loose stool; 1/2 caplet after each |
| children 9-11 years (60-95 lbs) | subsequent |
| | loose stool; but no more than 3 caplets in 24 hours |
| | 1 caplet after the first loose stool; 1/2 caplet after each |
| children 6-8 years (48-59 lbs) | subsequent |
| | loose stool; but no more than 2 caplets in 24 hours |
| children 2-5 years (34-47 lbs) | ask a doctor |
| children under 2 years (up to 33 | do not use |
| lbs) | uo not use |

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

exchange √ select[™]

Compare To The Active Ingredient of Imodium[®] A-D*

Anti-Diarrheal

LOPERAMIDE HCl 2 mg

(anti-diarrheal)

Controls the symptoms of diarrhea

Actual Size

12 CAPLETS

 $\sqrt{\text{quality value}}$

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D.

50844 REV0619M37502

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"

Manufactured For Your Military Exchanges Distributed by: LNK International, Inc., Hauppauge, NY 11788

1-800-426-9391

USB controls symptoms of diarrhea, including Travelers' Diarrhea Anti-diarmeal Loperamide HCI 2 mg Purpose Active ingredient (in each caplet) Drug Facts KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

NOT ink Jetted on .OT/EXP are embossed Questions or comments?

Drug Facts (continued)

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

exchange**Vselect**"

Compare To The Active Ingredient of Imodium® A-D*

Anti-Diarrheal LOPERAMIDE HCl 2 mg (anti-diarrheal)

Controls the symptoms of diarrhea

12 CAPLETS



Actual Size

quality value

AMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING . To open, tear

Use % to cut through plastic for caplet.

Johnson & Johnson Corporation, owner of the registerec trademark Imodium® A-D. 50844 REV0619M37502 *This product is not manufactured or distributed by

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"
Manufactured for Your Millary Exchanges
Distributed by: LINK International Inc., Hauppauge, NY 11788
1-800-426-8391

FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide Inactive ingredients corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate,

TOR NOR BROKEN ■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS Other information

Drug Facts (continued)

| esu fon ob | children under 2 years (up to 33 lbs) |
|--|---------------------------------------|
| ask a doctor | cal T4-45) years (34-47 lbs) |
| 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours | (sdl 63-84) sasey 8-8 inerblirb |
| 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours | (adl 39-09) arsay ff-9 nərblirk |
| 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours | adults and children 12 years and over |

■ find right dose on chart. If possible, use weight to dose; otherwise, use age. ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center If pregnant or breast-feeding, ask a health professional before use.

■ you get abdominal swelling or bulging. These may be signs of a serious condition. Stop use and ask a doctor if symptoms get worse ■ diarrhea lasts for more than 2 days

When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating

Ask a doctor or pharmacist before use it you are taking a prescription drug. Loperamide may interact with



Exchange Select 44-375

ANTI DIARRHEAL

loperamide hcl tablet

| P | rn | duct | Info | rmatio | 'n |
|---|----|------|------|----------|-----|
| _ | ıυ | uucı | THU | ı ına ul | ,,, |

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:55301-375

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| LOPERAMIDE HYDRO CHLO RIDE (UNII: 77TI35393C) (LOPERAMIDE - | LOPERAMIDE | 2 mg |
| UNII:6 X9O C3H4ID | HYDROCHLORIDE | 2 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULO SE, MICRO CRYSTALLINE (UNII: OP1R32D61U) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| SILICON DIO XIDE (UNII: ETJ7Z6 XBU4) | |
| DIBASIC CALCIUM PHO SPHATE DIHYDRATE (UNII: O7TSZ97GEP) | |

Product Characteristics

| Color | GREEN | Score | 2 pieces |
|----------|-------|--------------|----------|
| Shape | OVAL | Size | 10 mm |
| Flavor | | Imprint Code | 44;375 |
| Contains | | | |

Packaging

| F | ackaging | | | |
|---|------------------|--|-----------------------------|---------------------------|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:55301-375-02 | 2 in 1 CARTON | 05/03/2005 | |
| 1 | | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 2 | NDC:55301-375-22 | 1 in 1 CARTON | 05/03/2005 | |
| 2 | | 48 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| | | | | |

| Marketing Information | | | | | |
|-----------------------|--|----------------------|--------------------|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | |
| ANDA | ANDA076497 | 05/03/2005 | | | |
| | | | | | |

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

| Establishment | | | | |
|-------------------------|---------|-----------|----------------------------|--|
| Name | Address | ID/FEI | Business Operations | |
| LNK International, Inc. | | 038154464 | PACK(55301-375) | |

| Establishment | | | |
|-------------------------|---------|-----------|------------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 832867894 | MANUFACTURE(55301-375) |

| Establishment | | | | |
|-------------------------|---------|-----------|---------------------|--|
| Name | Address | ID/FEI | Business Operations | |
| LNK International, Inc. | | 967626305 | PACK(55301-375) | |

| Establishment | | | | |
|-------------------------|---------|-----------|---------------------|--|
| Name | Address | ID/FEI | Business Operations | |
| LNK International, Inc. | | 832867837 | PACK(55301-375) | |

| Establishment | | | |
|-------------------------|---------|-----------|---------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 868734088 | PACK(55301-375) |

Revised: 3/2020 ARMY AND AIR FORCE EXCHANGE SERVICE