

DAYTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated
Spirit Pharmaceuticals LLC

Daytime Severe Cold and Flu

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

• for the temporary relief of the following cold/flu symptoms: • minor aches and pains • headache • sore throat • nasal congestion • cough • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive • temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use

if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur • pain, nasal congestion or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed

adults and children 12 years & over	• take 2 caplets every 4 hours • swallow whole; do not crush, chew or dissolve • do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

store at 25°C (77°F); excursions permitted between 15°–30°C (59°–86°F)

Inactive Ingredients

croscarmellose sodium, D&C Yellow#10, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone*, pregelatinized starch, talc, titanium dioxide *may contain this ingredient

Questions or comments?

1-888-333-9792

Distributed by:

Cabinet Health P.B.C.

Principal Display panel



COMPOSTABLE
REFILL POUCH



CABINET:

Daytime Severe Cold & Flu Refill

COMPARE TO THE ACTIVE INGREDIENTS IN:

Tylenol® Cold + Flu Severe*

Acetaminophen 325 mg, Dextromethorphan HBr 10 mg,
Guaifenesin 200 mg, Phenylephrine HCl 5 mg

RELIEVES:

HEAD & BODY ACHES • FEVER & SORE THROAT • COUGH
NASAL CONGESTION • MUCUS & CHEST CONGESTION

30 CAPLETS



TO OPEN:

1. POSITION THUMBS INSIDE THE SHORT FLANGE OVER THE LOCKS.
2. GRIP WHILE PIVOTING THUMBS OUTWARD TO OPEN THE LOCKS.

Tamper evident: do not use if pouch is open

Drug Facts

Active ingredient (in each caplet)

Purpose

Acetaminophen 325 mg..... Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg..... Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Uses • for the temporary relief of the following cold/flu symptoms:
• minor aches and pains • headache • sore throat • nasal congestion
• cough • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive • temporarily reduces fever

Warnings **Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take:
• more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • If you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

• liver disease • heart disease • high blood pressure • thyroid disease
• diabetes • trouble urinating due to an enlarged prostate gland • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if • nervousness, dizziness, or sleeplessness occur • pain, nasal congestion or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warnings: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions • do not take more than directed

adults and children over 12 years old	• take 2 caplets every 4 hours • swallow whole; do not crush, chew or dissolve • do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information • store at 25°C (77°F); excursions permitted between 15°–30°C (59°–86°F)

Inactive ingredients Croscarmellose sodium, D&C Yellow #10, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone[†], pregelatinized starch, talc, titanium dioxide *may contain this ingredient

Questions or comments?

1-888-333-6792

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Cold + Flu Severe

† Less than the limit of detection and consistent with gluten-free diet labeling per FDA

NDC: 68210-4187-2

ACC-1-10033
Distributed by: Cabinet Health P.B.O.
Question/Comments? 1-888-333-6792
www.cabinethealth.com

Lot / Expiration ▼



Compostable
& Plastic Free



Independently
Quality Tested



Gluten Free
Verified[†]

DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride
tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68210-4221
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, POTATO (UNII: 8I089SAH3T)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	ET32
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-4221-3	30 in 1 JAR; Type 0: Not a Combination Product	11/14/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/14/2022	

Labeler - Spirit Pharmaceuticals LLC (179621011)

Revised: 12/2024

Spirit Pharmaceuticals LLC