

**YINCHIAO - acetaminophen and chlorpheniramine maleate tablet, film coated**  
**TIANJIN ZHONGXIN PHARMACEUTICAL GROUP CORP LTD LONGSHUNRONG**  
**PHARMACEUTICAL FACTORY**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**DRUG FACTS**

Active ingredient (in each tablet)

Acetaminophen 162.5 mg

Chlorpheniramine maleate 1 mg

Purpose

Pain reliever-fever reducer

Antihistamine

Uses

temporarily relieves these symptoms due to a cold, the flu, or hay fever:

minor aches and pains headache

sore throat

sneezing

runny nose

itchy, watery eyes

itching of the nose or throat

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

adult takes more than 24 tablets in 24 hours, which is the maximum daily amount for this product

child takes more than 5 doses in 24 hours

taken with other drugs containing acetaminophen

adult has 3 or more alcoholic drinks everyday while using this product

Do not give to children under 6 years of age unless directed by a doctor.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

whether a drug contains acetaminophen, ask a doctor or pharmacist.

for pain for more than 10 days (for adults) or 5 days (for children)

for fever for more than 3 days

Ask a doctor before use if the user has

liver disease

glaucoma

a breathing problem such as emphysema or chronic bronchitis

difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin

taking sedatives or tranquilizers

When using this product

do not exceed recommended dosage

avoid alcoholic beverages

may cause drowsiness (alcohol, sedatives, and tranquilizers may increase the drowsiness effect)

use caution when driving or operating machinery

may cause excitability especially in children

Stop use and ask a doctor if

sore throat is severe

sore throat persists for more than 2 days

sore throat is accompanied or followed by:

fever

headache

rash

nausea

vomiting

any of the following occurs (these could be signs of a serious condition):

pain or fever persists or gets worse or new symptoms occur

redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

adults and children 12 years of age and older: 4 tablets every 4 to 6 hours, not to exceed 6 doses (24 tablets) in 24 hours, or as directed by a doctor

children 6 to under 12 years of age: 2 tablets every 4 to 6 hours, not to exceed 5 doses (10 tablets) in 24 hours, or as directed by a doctor

children under 6 years: consult a doctor of age

#### Other information

keep tightly sealed

store between 15° to 30° C (59° to 86° F)

Questions or Comments? (888) 221-3496 M-F 9 am to 5 pm

you may also use this number to report serious adverse events associated with the use of this product

#### Inactive ingredients

Scutellaria baicalensis root, platycodon grandiflorum leaf, arctium lappa fruit, mint, FDC blue no. 2, FDC yellow no.5, FDC yellow no.6, titanium dioxide, soybean, forsythia suspensa fruit, lonicera japonica flower, hypromellose 2910 (15000 mpa.s), glycyrrhiza uralensis, notopterygium incisum root, phragmites australis root, polydextrose, polyethylene glycol 4000, schizonepeta tenifolia spike

YINCHIAO, NDC58975-022-01, Acetaminophen - Pain Reliever-Fever Reducer, Chlorpheniramine Maleate - Antihistamine, See new warnings information, 120 Tablets



## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58975-022
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	162.5 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg

## Inactive Ingredients

Ingredient Name	Strength
SCUTELLARIA BAICALENSIS ROOT (UNII: 7J95K7ID2S)	
PLATYCODON GRANDIFLORUM LEAF (UNII: 2L64H2X8DY)	
ARCTIUM LAPPA FRUIT (UNII: EA541308MV)	
MINT (UNII: FV98Z8GITP)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SOYBEAN (UNII: L7HT8F1ZOD)	
FORSYTHIA SUSPENS A FRUIT (UNII: P4793M1ES5)	
LONICERA JAPONICA FLOWER (UNII: 4465L2WS4Y)	
HYPROMELLOSE 2910 (15000 MPA.S) (UNII: 288VBX44JC)	
GLYCYRRHIZA URALENSIS (UNII: 42B5YD8F0K)	
NOTOPTERYGIUM INCISUM ROOT (UNII: 5Z2WW4J6RI)	
PHRAGMITES AUSTRALIS ROOT (UNII: BBJ7AOM815)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HF16D95)	
SCHIZONEPETA TENUFOLIA SPIKE (UNII: 2FN3BA1MZE)	

## Product Characteristics

<b>Color</b>	yellow (YC)	<b>Score</b>	no score
<b>Shape</b>	ROUND (YC)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	YC
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58975-022-01	120 in 1 BOX		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC monograph final	part341	02/16/2011	
---------------------	---------	------------	--

**Labeler** - TIANJIN ZHONGXIN PHARMACEUTICAL GROUP CORP LTD LONGSHUNRONG PHARMACEUTICAL FACTORY (654133313)

Revised: 2/2011

TIANJIN ZHONGXIN PHARMACEUTICAL GROUP CORP LTD LONGSHUNRONG PHARMACEUTICAL FACTORY