# AMPROMAX- amprolium oral solution solution Vetr LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

#### Vetr Ampromax (amprolium, 9.6% oral solution)

WARNING: NOT FOR USE IN HUMANS. Keep this and all drugs out of the reach of children. FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with water and seek medical attention. RESUDUE WARNING: Withdraw 24 h before slaughter. A withdrawl period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

#### Shake well before use.

<u>5 Day Regimen</u>(coccidiosis symptoms present): Add Ampromax (9.6% amprolium) oral solution to drinking water at the rate of 16 fl oz/100 gal (473 mL/378.5 mL). At the usual rate of water consumption, this will provide intake of approximately 10 mg amprolium/kg body weight (2.2 lbs). Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed and all calves should be treated.

<u>21 Day Regimen</u>(coccidiosis exposure risk): During periods of exposure, or when experience indicates environmental coccidiosis risk, **add Ampromax (9.6% amprolium) oral solution to drinking water at the rate of 8 fl oz/100 gal** (237 mL/378.5 mL). At usual rates of consumption, this will provide approximately 5 mg amprolium/kg body weight (2.2 lbs). Offer this solution as the only source of water for 21 days.

Store at room temperature, at or below 25 C (<77 F)

Amprolium USP, 96 mg/mL (9.6% w/v)

Purified water, preservative.

# AmproMax

## Amprolium 9 6mg/mL

(Shake Well)

INDICATIONS: Coccidiosis caused by Eimeria bovis and E zuernii in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before

WARNING: NOT FOR USE IN HUMANS. Keep this and all

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY, MAY

**RESIDUE WARNINGS:** Withdraw 24 hours before

#### 500 mL/16.9oz

# **AmproMax** 9.6% Oral Solution

(Shake Well)

**INDICATIONS:** Coccidiosis caused by Eimeria bovis and E. zuernii in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

WARNING: NOT FOR USE IN HUMANS. Keep this and all drugs out of the reach of children.

**PRECAUTIONS:** FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

**RESIDUE WARNINGS:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

## **One Gallon**

#### Active Constituents:

Amprolium USP 96 mg/mL (9.6%) Other Ingredients: Water

#### Directions & Storage

#### 5 Day Regimen

(When Coccidiosis symptoms are present): Add AMPROLIUM 9.6% Oral Solution to drinking water at

the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.



AM001

#### 21 Day Regimen (When Coccidiosis exposure is a risk):

During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AmproMax 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of 02772 00409 7 water for 21 days.

KEEP OUT OF THE REACH OF CHILDREN

#### **Active Constituents:**

Amprolium USP 96 mg/mL (9.6%) Other Ingredients: Water

#### **Directions & Storage**

#### 5 Day Regimen

#### (When Coccidiosis symptoms are present):

Add AMPROLIUM 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.



AM002

#### 21 Day Regimen (When Coccidiosis exposure is a risk):

During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add AmproMax 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of 6 02772 00410 3 water for 21 days.

KEEP OUT OF THE REACH OF CHILDREN

## AMPROMAX

amprolium oral solution solution

Product Information								
Product Type		OTC ANIMAL DRUG	Item Coo	tem Code (Source)		NDC:86213-772		
Route of Admin	nistration	ORAL						
Active Ingredient/Active Moiety								
Ingredient Name				Basis of Strength Strength				
AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307K				AMPROLIUM 96 mg in 1 m		96 mg in 1 mL		
Inactive Ingredients								
Ingredient Name					Strength			
WATER (UNII: 059QF0KO0R)								
Packaging								
Packaging # Item Cod	e Packa	age Description	Marketin	g Start Da	ite Market	ing End Date		
		age Description 1 BOTTLE, PLASTIC	Marketin	g Start Da	ite Market	ing End Date		
# Item Cod	2-01 500 mL in	• •	Marketin	g Start Da	ite Market	ing End Date		
# Item Cod 1 NDC:86213-772	2-01 500 mL in	1 BOTTLE, PLASTIC	Marketin	g Start Da	ite Market	ing End Date		
# Item Cod 1 NDC:86213-772	2-01 500 mL in 2 2-02 3785.4 mL	1 BOTTLE, PLASTIC in 1 BOTTLE, PLASTIC	Marketin	g Start Da	ite Market	ing End Date		
<ul> <li># Item Cod</li> <li>1 NDC:86213-772</li> <li>2 NDC:86213-772</li> </ul>	2-01 500 mL in 2-02 3785.4 mL	1 BOTTLE, PLASTIC in 1 BOTTLE, PLASTIC		g Start Da Marketing Dat	g Start M	ing End Date arketing End Date		
<ul> <li># Item Cod</li> <li>1 NDC:86213-772</li> <li>2 NDC:86213-772</li> <li>Marketing</li> <li>Marketing</li> </ul>	2-01 500 mL in 2-02 3785.4 mL	I BOTTLE, PLASTIC in 1 BOTTLE, PLASTIC ion tion Number or Mono	graph	Marketing	g Start M	arketing End		

Labeler - Vetr LLC (132219984)

Establishment						
Name	Address	ID/FEI	<b>Business Operations</b>			
Vetr LLC		132219984	manufacture, label			

Revised: 6/2025

Vetr LLC