

**CITROMA- magnesium citrate liquid**  
**Western Family Foods, Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active Ingredient**

Magnesium citrate 1.745 g per fl oz

**Purpose**

Saline Laxative

**Use**

for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

**Warnings**

**Do not use**

- if you are on a low salt diet

**Ask a doctor**

**before use if you have**

- kidney disease
- a magnesium or sodium-restricted diet
- abdominal pain, nausea or vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already use a laxative for a period longer than 1 week

**Ask a doctor or pharmacist before use if you are**

taking any other drugs. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop Use and ask a doctor if**

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age - ask a doctor

discard unused product within 24 hours of opening bottle

**Other information***Other information*

- magnesium content 290 mg per 1 fl oz (30 mL)
- sodium content 45 mg per 1 fl oz (30 mL)
- store at temperatures between 46° and 86° F (8° and 30° C)

**Inactive ingredients**

citric acid, lemon oil, polyethylene glycol, purified water, sodium bicarbonate, sodium saccharin, sucrose

**Questions or comments**

1-800-525-2558

**Adverse Reactions**

Proudly Distributed by Western Family Foods, Inc.

P.O.Box 4057, Portland, OR 97208 U.S.A.

Copyright 2014 Marca Registrada

[www.westernfamily.com](http://www.westernfamily.com)

**Principal Display Panel**

WESTERN

FAMILY

Magnesium

Citrate


Saline Laxative

Oral Solution

**TAMPER EVIDENT:**

DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.


10 FL OZ (296 mL)



# Magnesium Citrate

Saline Laxative Oral Solution

**TAMPER EVIDENT:**  
DO NOT USE IF THE  
BREAKAWAY BAND  
ON CAP IS BROKEN  
OR MISSING.



**10 FL OZ (296 mL) lemon**

| Drug Facts  |  |
|---|--|
| <b>Active ingredient</b><br>Magnesium citrate<br>1.745 g per fl oz ..... Saline laxative  | <b>Purpose</b><br>Saline laxative            |
| <b>Use</b> for relief of occasional constipation (irregularity). Generally produces bowel movement in ½ to 6 hours  |  |
| <b>Warnings</b><br><b>Do not use</b><br>■ if you are on a low salt diet<br><b>Ask a doctor before use if you have</b><br>■ kidney disease ■ a magnesium or sodium-restricted diet<br>■ abdominal pain, nausea, or vomiting<br>■ noticed a sudden change in bowel habits that persists over a period of 2 weeks<br>■ already used a laxative for a period longer than 1 week<br><b>Ask a doctor or pharmacist before use if you are</b><br>taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.<br><b>Stop use and ask a doctor if you have</b> rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.<br><b>If pregnant or breast-feeding,</b> ask a health professional before use.<br><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a |  |
| Drug Facts (continued)  |  |
| Poison Control Center right away.   |  |
| <b>Directions</b> ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses  |  |
| adults and children 12 years of age and over  | 6.5 to 10 fl oz maximum 10 fl oz in 24 hours |
| children 6 to under 12 years of age   | 3 to 7 fl oz maximum 7 fl oz in 24 hours     |
| children 2 to under 6 years of age  | 2 to 3 fl oz maximum 3 fl oz in 24 hours     |
| children under 2 years of age   | ask a doctor                                 |
| discard unused product within 24 hours of opening bottle  |  |
| <b>Other information</b><br>■ magnesium content 290 mg per 1 fl oz (30 mL)<br>■ sodium content 45 mg per 1 fl oz (30 mL)<br>■ store at temperatures between 46° and 86° F (8° and 30° C)  |  |
| <b>Inactive ingredients</b> citric acid, lemon oil, polyethylene glycol, purified water, sodium bicarbonate, sodium saccharin, sucrose  |  |
| <b>Questions? Call 1-800-525-2558</b>   |  |

UPC 0-15400-03426-5

80MAG 00000

QUALITY

Proudly Distributed by Western Family Foods, Inc.  
P.O. Box 4057, Portland, OR 97208 U.S.A.  
Copyright © 2014 Marca Registrada  
www.westernfamily.com 03426-A-VI L0003073SB

## CITROMA

magnesium citrate liquid

### Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:55312-667 |
| Route of Administration | ORAL           |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength            |
|---|-------------------|---------------------|
| MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838) | MAGNESIUM CITRATE | 1.745 g in 29.61 mL |

### Inactive Ingredients

| Ingredient Name                            | Strength |
|--|----------|
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) |          |
| LEMON OIL (UNII: I9GRO824LL)               |          |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)    |          |
| WATER (UNII: 059QF0KO0R)                   |          |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO)      |          |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY)        |          |
| SUCROSE (UNII: C151H8M554)                 |          |

**Packaging**

| # | Item Code        | Package Description       | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------------|----------------------|--------------------|
| 1 | NDC:55312-667-38 | 296 mL in 1 BOTTLE, GLASS |                      |                    |

**Marketing Information**

| Marketing Category      | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part334                                  | 04/15/1990           |                    |

**Labeler** - Western Family Foods, Inc (192166072)**Registrant** - Vi-Jon (790752542)**Establishment**

| Name   | Address | ID/FEI    | Business Operations    |
|--------|---------|-----------|------------------------|
| Vi-Jon |         | 790752542 | manufacture(55312-667) |

Revised: 11/2014

Western Family Foods, Inc