

**LANSOPRAZOLE- lansoprazole capsule, delayed release**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Lansoprazole Delayed-Release Capsules 15 mg Drug Facts**

**Active ingredient (in each capsule)**

Lansoprazole 15 mg

**Purpose**

Acid Reducer

**Use**

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:** Do not use if you are allergic to lansoprazole

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are**  
taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medicine)

### **Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

### **14-Day Course of Treatment**

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

### **Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

### **Other information**

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

## **Inactive ingredients**

D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, gelatin, hypromellose, low substituted hydroxypropyl cellulose, mannitol, meglumine, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

## **Questions or comments?**

**1-800-719-9260**

## **Consumer Information**

### ***Treats Frequent Heartburn***

### ***Lansoprazole***

### ***Delayed-Release Capsules 15 mg / Acid Reducer***

- May take 1 to 4 days for full effect
- Sodium Free

**Please read the entire package insert before taking Lansoprazole Delayed-Release Capsules 15 mg.**

**Save for future reference.**

### **How Lansoprazole Delayed-Release Capsules 15 mg Treats Your Frequent Heartburn**

Lansoprazole Delayed-Release Capsules 15 mg stops acid production at the source – the **pumps** that release acid into the stomach. Lansoprazole Delayed-Release Capsules 15 mg is taken once a day (every 24 hours), every day for 14 days.

### **What You Can Expect When Taking Lansoprazole Delayed-Release Capsules 15 mg**

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take Lansoprazole Delayed-Release Capsules 15 mg in the morning before eating. Lansoprazole Delayed-Release Capsules 15 mg is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take Lansoprazole Delayed-Release Capsules 15 mg every day for 14 days to treat your frequent heartburn.

### **Who Should Take Lansoprazole Delayed-Release Capsules 15 mg**

Adults (18 years and older) with **frequent heartburn** – when you have heartburn 2 or more days a week.

### **Who Should NOT Take Lansoprazole Delayed-Release Capsules 15 mg**

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

### **How to Take Lansoprazole Delayed-Release Capsules 15 mg**

#### **14-DAY Course of Treatment**

- Swallow 1 capsule with a glass of water before eating in the morning.
- Take every day for 14 days.
- Do not take more than 1 capsule a day.
- Swallow whole. Do not crush or chew capsules.
- Do not use for more than 14 days unless directed by your doctor.

### **When to Take Lansoprazole Delayed-Release Capsules 15 mg Again**

**You may repeat a 14-day course of therapy every 4 months.**

## **When to Talk to Your Doctor**

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

## **Warnings and When to Ask Your Doctor**

**Allergy alert:** Do not use if you are allergic to lansoprazole

### **Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

### **Ask a doctor before use if you have**

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

### **Ask a doctor or pharmacist before use if you are taking**

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medicine)

### **Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.

- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

### **Clinical studies prove Lansoprazole Delayed-Release Capsules 15 mg effectively treats frequent heartburn**

In three clinical studies, Lansoprazole Delayed-Release Capsules 15 mg was shown to be significantly better than placebo in treating frequent heartburn.

### **How Lansoprazole Delayed-Release Capsules 15 mg is Sold**

Lansoprazole Delayed-Release Capsules 15 mg is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

### **For Questions or Comments About Lansoprazole Delayed-Release Capsules 15 mg**

**Call 1-800-719-9260**

Distributed By

Perrigo®

Allegan, MI 49010

: 3T300 00 J5

### **Package/Label Principal Display Panel**

Compare to Prevacid® 24 HR

active ingredient

24 HOUR

lansoprazole

Delayed Release Capsules, 15 mg

Acid Reducer

TREATS FREQUENT HEARTBURN

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

SODIUM FREE

14 Capsules

one 14-day course of treatment

actual size



**meijer**

NDC 41250-117-01

Compare to Prevacid® 24 HR active ingredient\*

**24 HOUR**

**lansoprazole**

Delayed Release Capsules, 15 mg  
Acid Reducer


**TREATS FREQUENT HEARTBURN**

MAY TAKE  
1 TO 4 DAYS  
FOR FULL EFFECT  
SODIUM FREE

**14 Capsules**  
one 14-day course of treatment



actual size  
1 bottle inside



**TREATS FREQUENT HEARTBURN**

Delayed Release Capsules, 15 mg  
Acid Reducer

**lansoprazole**

**24 HOUR**

**meijer**

**Drug Facts**

**Active ingredient (in each capsule)** Purpose  
Lansoprazole 15 mg Acid Reducer

**Use**

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:** Do not use if you are allergic to lansoprazole. Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness

**Drug Facts (continued)**

- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days

- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

**Drug Facts (continued)**

**Other information**

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

**Inactive ingredients**

D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, gelatin, hypromellose, low substituted hydroxypropyl cellulose, mannitol, magnesium, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
 ■ frequent chest pain  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss  
 ■ nausea or vomiting ■ stomach pain  
 Ask a doctor or pharmacist before use if you are taking  
 ■ warfarin (blood-thinning medicine)  
 ■ prescription antifungal or anti-yeast medicines  
 ■ digoxin (heart medicine) ■ theophylline (asthma medicine)  
 ■ tacrolimus or mycophenolate mofetil (immune system medicine)

■ swallow 1 capsule with a glass of water before eating in the morning  
 ■ take every day for 14 days  
 ■ do not take more than 1 capsule a day  
 ■ swallow whole. Do not crush or chew capsules.  
 ■ do not use for more than 14 days unless directed by your doctor  
**Repeated 14-Day Courses (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor  
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Questions or comments?**  
 1-800-719-9280

**KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.**

**DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING OR WHITE BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.**

DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, MI 49544  
 www.meijer.com



PD 3066414



\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Protonix®.

3T3D6 6E C6

## LANSOPRAZOLE

lansoprazole capsule, delayed release

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-117
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LANSOPRAZOLE (UNII: 0K5C5T2QPG) (LANSOPRAZOLE - UNII:0K5C5T2QPG)	LANSOPRAZOLE	15 mg

### Inactive Ingredients

Ingredient Name	Strength
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
HYDROXYPROPYL CELLULOSE, LOW SUBSTITUTED (UNII: 2165RE0K14)	
MANNITOL (UNII: 3OWL53L36A)	
MEGLUMINE (UNII: 6HG8UB2MUY)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	PINK, GREEN	Score	no score
-------	-------------	-------	----------

<b>Shape</b>	CAPSULE	<b>Size</b>	16 mm
<b>Flavor</b>		<b>Imprint Code</b>	L3T3
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-117-01	1 in 1 CARTON	05/18/2012	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-117-02	2 in 1 CARTON	05/18/2012	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-117-03	3 in 1 CARTON	05/18/2012	
3		14 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202319	05/18/2012	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 12/2019

Meijer Distribution Inc