

**TEXACLEAR KIDS DAYTIME COUGH AND CHEST CONGESTION-  
dextromethorphan hbr, guaifenesin liquid  
GM Pharmaceuticals, INC**

-----  
**TexaClear Kids Daytime Cough and Chest Congestion**

**TexaClear Kids Daytime Cough and Chest Congestion  
(dextromethorphan HBr, guaifenesin), solution  
GM Pharmaceuticals, Inc.**

**Drug Facts**

***Active ingredients (in each 15 mL)***

Dextromethorphan HBr 5 mg  
Guaifenesin 100 mg

***Purpose***

Cough suppressant  
Expectorant

***Uses***

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.
- temporarily relieves:
  - the intensity of coughing
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

**Do not exceed recommended dosage.**

**Do not use**

in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- persistent or chronic cough such as occurs with asthma
- a cough that occurs with too much phlegm (mucus)
- a sodium-restricted diet

**Stop use and ask a doctor if**

cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not exceed 6 doses per 24-hour period.
- measure only with the dosing cup provided.
- mL= milliliter

<b>Age</b>	<b>Dose</b>
children 6 years to under 12 years	30 mL (2 dose cups) every 4 hours
children 4 years to under 6 years	15 mL (1 dose cup) every 4 hours
<b>children under 4 years</b>	<b>do not use.</b>

**Other information**

- each 15 mL contains: **Sodium 7.5 mg**
- store at room temperature 20-30°C (68-86°F)

**Inactive ingredients**

citric acid, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol, sucralose

**Questions or comments?**

Call 1-888-535-0305 9 a.m. - 5 p.m. CST.

**Distributed by:**

GM Pharmaceuticals, Inc.  
Fort Worth, TX. 76118

**Principal Display Panel**

**TexaClear Kids Daytime Cough and Chest Congestion**

NDC 58809-216-08

8 fl. oz. (237 mL)

Dextromethorphan HBR – Cough Suppressant  
Guaifenesin – Expectorant

Tamper evident: do not use if foil seal under cap is broken on missing

- Dye Free
- Sugar Free
- Gluten Free

- Alcohol Free
- Acetaminophen Free

Ages 4+

Relieves:

- Chest Congestion
- Thins & Loosens Mucus
- Cough

Distributed by:

GM Pharmaceuticals, Inc. Fort Worth, TX 76118

NDC 58809-216-08

**TEXACLEAR**

**Kids**

**DAYTIME  
COUGH &  
CHEST CONGESTION**

Dextromethorphan HBr 5 mg- Cough Suppressant  
Guaifenesin 100 mg- Expectorant

**FREE & CLEAR  
FORMULA**

Chest Congestion  
Thins & Loosens Mucus  
Cough

8 fl oz (237 mL)

**LIQUID FAST**

**FREE & CLEAR  
CLEAN FORMULA**

- Dye Free
- Sugar Free
- Gluten Free
- Alcohol Free
- Acetaminophen Free

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP IS BROKEN OR MISSING

LIFT HERE FOR COMPLETE DRUG FACTS

- Ages 4+
- Non-drowsy

Please visit our website  
[www.texaclear.com](http://www.texaclear.com)

3 58809 21608 6

Distributed by:  
GM Pharmaceuticals, Inc.  
Fort Worth, TX 76118

401148-01 Rev. 0923

# Drug Facts

## Active ingredients (in each 15 mL)

Dextromethorphan HBr 5 mg.....Cough suppressant  
Guaifenesin 100 mg .....Expectorant

## Purpose

## Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.
- temporarily relieves:
  - the intensity of coughing
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

## Warnings

**Do not exceed recommended dosage.**

**Do not use** in a child who is taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- a sodium-restricted diet

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- do not exceed 6 doses per 24-hour period
- measure only with dosing cup provided
- mL = milliliter

Age	Dose
children 6 years to under 12 years	30 mL (2 dose cups) every 4 hours
children 4 years to under 6 years	15 mL (1 dose cup) every 4 hours
<b>children under 4 years</b>	<b>do not use</b>

## Other information

- each 15 mL contains: **Sodium 7.5 mg**
- store at room temperature 20-30°C (68-86°F)

## Inactive ingredients

citric acid, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol, sucralose

**Questions or comments?** Call 1-888-535-0305

## TEXACLEAR KIDS DAYTIME COUGH AND CHEST CONGESTION

dextromethorphan hbr, guaifenesin liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58809-216
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 15 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58809-216-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/25/2023	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/25/2023	

**Labeler** - GM Pharmaceuticals, INC (793000860)