

COLD AND FLU HBP- acetaminophen, chlorpheniramine maleate tablet, film coated
WALMART INC.

Equate 44-692

Active ingredients (in each tablet)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg

Purpose

Pain reliever/fever reducer
Antihistamine

Uses

- temporarily relieves:
 - headache
 - sneezing
 - runny nose
 - minor aches and pains
 - sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if the user is

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days (children 6 to under 12 years) or 10 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

adults and children 12 years and over	2 tablets every 4 to 6 hours; do not take more than 12 tablets in 24 hours
children 6 to under 12	1 tablet every 4 to 6 hours; do not take more than 5

years	tablets in 24 hours
children under 6 years	do not use unless directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-888-287-1915

Principal display panel

equate[™]

NDC 79903-120-09

Compare to
Coricidin® HBP
Cold & Flu
active
ingredients*

DECONGESTANT-FREE

Cold & Flu HBP

Acetaminophen - Pain reliever/Fever reducer
Chlorpheniramine maleate - Antihistamine

FOR PEOPLE WITH HIGH BLOOD PRESSURE

Relieves:

- Runny nose
- Sneezing
- Body aches & pains
- Fever, sore throat

Actual Size

**20
TABLETS**

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Satisfaction guaranteed – For questions or comments please call **1-888-287-1915**.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Cold & Flu.
50844 REV1020A69209

Drug Facts KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine

Drug Facts (continued)

Uses

- temporarily relieves:
 - headache
 - sneezing
 - runny nose
 - minor aches and pains
 - sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if the user is

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness

Drug Facts (continued)

see end flap for expiration date and lot number

Drug Facts (continued)

Warnings

- drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children
- avoid alcoholic beverages

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days (children 6 to under 12 years) or 10 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed

adults and children 12 years and over	2 tablets every 4 to 6 hours; do not take more than 12 tablets in 24 hours
children 6 to under 12 years	1 tablet every 4 to 6 hours; do not take more than 5 tablets in 24 hours
children under 6 years	do not use unless directed by a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Drug Facts (continued)

Inactive ingredients corn starch, croscollon, FDCC red #40, aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, potassium silicate, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-888-287-1915

equate™

DECONGESTANT-FREE

Cold & Flu HBP

Acetaminophen - Pain reliever/Fever reducer
Chlorpheniramine maleate - Antihistamine

FOR PEOPLE WITH HIGH BLOOD PRESSURE

Relieves:

- Runny nose
- Sneezing
- Body aches & pains
- Fever, sore throat

20 TABLETS

Actual Size

COLD AND FLU HBP			
acetaminophen, chlorpheniramine maleate tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-120

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	19mm
Flavor		Imprint Code	44;692
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-120-09	2 in 1 CARTON	04/14/2022	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/14/2022	

Labeler - WALMART INC. (051957769)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(79903-120) , pack(79903-120)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(79903-120)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(79903-120)

Revised: 6/2025

WALMART INC.