


BACLOFEN- baclofen tablet
Lifestar Pharma LLC

Baclofen Tablets, USP
Rx only

DESCRIPTION

Baclofen, USP is a muscle relaxant and antispastic.

Its chemical name is 4-amino-3-(4-chlorophenyl)-butanoic acid. The structural formula is:



$\text{C}_{10}\text{H}_{12}\text{ClNO}_2$ M.W. 213.66

Baclofen, USP is a white to off-white odorless or practically odorless crystalline powder. It is slightly soluble in water, very slightly soluble in methanol and insoluble in chloroform.

Each tablet, for oral administration, contains 5 mg, 10 mg or 20 mg baclofen, USP. In addition, each tablet contains the following inactive ingredients: anhydrous lactose, colloidal silicon dioxide, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose and sodium starch glycolate.

CLINICAL PHARMACOLOGY

The precise mechanism of action of baclofen is not fully known. Baclofen is capable of inhibiting both monosynaptic and polysynaptic reflexes at the spinal level, possibly by hyperpolarization of afferent terminals, although actions at supraspinal sites may also occur and contribute to its clinical effect. Although baclofen is an analog of the putative inhibitory neurotransmitter gamma-aminobutyric acid (GABA), there is no conclusive evidence that actions on GABA systems are involved in the production of its clinical effects. In studies with animals baclofen has been shown to have general CNS depressant properties as indicated by the production of sedation with tolerance, somnolence, ataxia, and respiratory and cardiovascular depression. Baclofen is rapidly and extensively absorbed and eliminated. Absorption may be dose-dependent, being reduced with increasing doses. Baclofen is excreted primarily by the kidney in unchanged form and there is relatively large intersubject variation in absorption and/or elimination.

INDICATIONS AND USAGE

Baclofen tablets are useful for the alleviation of signs and symptoms of spasticity resulting from multiple sclerosis, particularly for the relief of flexor spasms and concomitant pain, clonus, and muscular rigidity.

Patients should have reversible spasticity so that baclofen treatment will aid in restoring residual function. Baclofen tablets may also be of some value in patients with spinal cord injuries and other spinal cord diseases.

Baclofen tablets are not indicated in the treatment of skeletal muscle spasm resulting from rheumatic disorders.

The efficacy of baclofen in stroke, cerebral palsy, and Parkinson's disease has not been established and, therefore, it is not recommended for these conditions.

CONTRAINDICATIONS

Hypersensitivity to baclofen.

WARNINGS

a. Neonatal Withdrawal Symptoms: Withdrawal symptoms have been reported starting hours to days after delivery in neonates whose mothers were treated with oral baclofen throughout pregnancy. The symptoms of withdrawal in these infants have included increased muscle tone, tremor, jitteriness, and seizure. If the potential benefit justifies the potential risk to the fetus and oral baclofen is continued during pregnancy, gradually reduce the dose and discontinue baclofen before delivery. If slow withdrawal is not feasible, advise the parents or caregivers of the potential for neonatal withdrawal.

b. Abrupt Drug Withdrawal: Hallucinations and seizures have occurred on abrupt withdrawal of baclofen. Therefore, except for serious adverse reactions, the dose should be reduced slowly when the drug is discontinued.

c. Impaired Renal Function: Because baclofen is primarily excreted unchanged through the kidneys, it should be given with caution, and it may be necessary to reduce the dosage.

d. Stroke: Baclofen has not significantly benefited patients with stroke. These patients have also shown poor tolerability to the drug.

e. Pregnancy: Baclofen has been shown to increase the incidence of omphaloceles (ventral hernias) in fetuses of rats given approximately 13 times the maximum dose recommended for human use, at a dose which caused significant reductions in food intake and weight gain in dams. This abnormality was not seen in mice or rabbits.

There was also an increased incidence of incomplete sternebral ossification in fetuses of rats given approximately 13 times the maximum recommended human dose, and an increased incidence of unossified phalangeal nuclei of forelimbs and hindlimbs in fetuses of rabbits given approximately 7 times the maximum recommended human dose. In mice, no teratogenic effects were observed, although reductions in mean fetal weight with consequent delays in skeletal ossification were present when dams were given 17

and 34 times the human daily dose. There are no studies in pregnant women. Baclofen should be used during pregnancy only if the benefit clearly justifies the potential risk to the fetus.

PRECAUTIONS

Because of the possibility of sedation, patients should be cautioned regarding the operation of automobiles or other dangerous machinery, and activities made hazardous by decreased alertness. Patients should also be cautioned that the central nervous system effects of baclofen may be additive to those of alcohol and other CNS depressants.

Baclofen should be used with caution where spasticity is utilized to sustain upright posture and balance in locomotion or whenever spasticity is utilized to obtain increased function.

In patients with epilepsy, the clinical state and electroencephalogram should be monitored at regular intervals, since deterioration in seizure control and EEG have been reported occasionally in patients taking baclofen.

It is not known whether this drug is excreted in human milk. As a general rule, nursing should not be undertaken while a patient is on a drug since many drugs are excreted in human milk.

A dose-related increase in incidence of ovarian cysts and a less marked increase in enlarged and/or hemorrhagic adrenal glands was observed in female rats treated chronically with baclofen.

Ovarian cysts have been found by palpation in about 4% of the multiple sclerosis patients that were treated with baclofen for up to one year. In most cases these cysts disappeared spontaneously while patients continued to receive the drug. Ovarian cysts are estimated to occur spontaneously in approximately 1% to 5% of the normal female population.

Pediatric Use

Safety and effectiveness in pediatric patients below the age of 12 years have not been established.

ADVERSE REACTIONS

The most common is transient drowsiness (10 to 63%). In one controlled study of 175 patients, transient drowsiness was observed in 63% of those receiving baclofen compared to 36% of those in the placebo group. Other common adverse reactions are dizziness (5 to 15%), weakness (5 to 15%) and fatigue (2 to 4%).

Others reported:

Neuropsychiatric: Confusion (1 to 11%), headache (4 to 8%), insomnia (2 to 7%); and, rarely, euphoria, excitement, depression, hallucinations, paresthesia, muscle pain, tinnitus, slurred speech, coordination disorder, tremor, rigidity, dystonia, ataxia, blurred vision, nystagmus, strabismus, miosis, mydriasis, diplopia, dysarthria, epileptic seizure.

Cardiovascular: Hypotension (0 to 9%). Rare instances of dyspnea, palpitation, chest pain, syncope. Gastrointestinal: Nausea (4 to 12%), constipation (2 to 6%); and rarely, dry mouth, anorexia, taste disorder, abdominal pain, vomiting, diarrhea, and positive test for occult blood in stool.

Genitourinary: Urinary frequency (2 to 6%); and rarely, enuresis, urinary retention, dysuria, impotence, inability to ejaculate, nocturia, hematuria.

Other: Instances of rash, pruritus, ankle edema, excessive perspiration, weight gain, nasal congestion. Some of the CNS and genitourinary symptoms may be related to the underlying disease rather than to drug therapy. The following laboratory tests have been found to be abnormal in a few patients receiving baclofen: increased SGOT, elevated alkaline phosphatase, and elevation of blood sugar.

To report SUSPECTED ADVERSE REACTIONS, contact Lifestar Pharma LLC at 1-888-995-4337 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Signs and Symptoms: Vomiting, muscular hypotonia, drowsiness, accommodation disorders, coma, respiratory depression and seizures.

Treatment: In the alert patient, empty the stomach promptly by induced emesis followed by lavage. In the obtunded patient, secure the airway with a cuffed endotracheal tube before beginning lavage (do not induce emesis). Maintain adequate respiratory exchange, do not use respiratory stimulants.

DOSAGE AND ADMINISTRATION

The determination of optimal dosage requires individual titration. Start therapy at a low dosage and increase gradually until optimum effect is achieved (usually between 40 to 80 mg daily).

The following dosage titration schedule is suggested:

5 mg t.i.d. for 3 days

10 mg t.i.d. for 3 days

15 mg t.i.d. for 3 days

20 mg t.i.d. for 3 days

Thereafter additional increases may be necessary but the total daily dose should not exceed a maximum of 80 mg daily (20 mg q.i.d.).

The lowest dose compatible with an optimal response is recommended. If benefits are not evident after a reasonable trial period, patients should be slowly withdrawn from the drug (see **WARNINGS**, [Abrupt Drug Withdrawal](#)).

HOW SUPPLIED

Baclofen Tablets USP, 5 mg are available as a white to off white, round, flat-faced, beveled edge tablet, debossed with "LS" on one side and "85" on the other side,

containing 5 mg baclofen, USP are supplied as follows:

Bottle of 100 tablets, with child-resistant closure. NDC 70756-085-11

Baclofen Tablets USP, 10 mg are available as a white to off white, round, flat-faced, beveled edge tablet, debossed with "LS 288" on one side and functional scored on the other side, containing 10 mg baclofen, USP are supplied as follows:

Bottle of 100 tablets, with child-resistant closure. NDC 70756-288-11

Bottle of 1000 tablets. This package is not child resistant. NDC 70756-288-12

Unit-dose child-resistant blister pack of 100 (10 x 10) tablets. NDC 70756-288-99

NDC 70756-288-12 in bottle of 1000 tablets. This package is not child resistant.

Baclofen Tablets USP, 20 mg are available as a white to off white, round, flat-faced, beveled edge tablet debossed with "LS 289" on one side and functional scored on the other side, containing 20 mg baclofen, USP are supplied as follows:

Bottle of 100 tablets, with child-resistant closure. NDC 70756-289-11

Bottle of 500 tablets. This package is not child resistant. NDC 70756-289-51

Bottle of 1000 tablets. This package is not child resistant. NDC 70756-289-12

Unit-dose child-resistant blister pack of 100 (10 x 10) tablets. NDC 70756-289-99

Dispense in a well-closed container as defined in the USP, with a child-resistant closure.

Store at 20° to 25°C [68° to 77°F] [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for:

Lifestar Pharma LLC

1200 MacArthur Blvd.

Mahwah, NJ 07430 USA

Product of Poland

or

Manufactured for:

Lifestar Pharma LLC

1200 MacArthur Blvd.

Mahwah, NJ 07430 USA

Made in India

Revised: October 2023, V-05

NDC 70756-085-11

Baclofen


Tablets, USP

5 mg

Rx only

100 TABLETS

Product of Poland

<p>NDC 70756-085-11</p> <p>Baclofen Tablets, USP</p> <p>5 mg</p> <p>Rx only 100 Tablets</p> <p> Lifestar Serving Life</p>	<p>Each tablet contains Baclofen, USP 5 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194 ECYYTLJKJKUS176600 Rev: 12/21,V-02</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Product of Poland</p> <p>12004958</p> <p> GTIN SN EXP LOT</p> <p> 10129</p> <p> 317075610851109</p> <p>45x15 mm Unvarnished area</p>
--	--	--

NDC 70756-085-11

Baclofen


Tablets, USP

5 mg

Rx only

100 TABLETS

Made in India

<p>NDC 70756-085-11</p> <p>Baclofen Tablets, USP</p> <p>5 mg</p> <p>Rx only 100 Tablets</p> 	<p>Each tablet contains Baclofen, USP 5 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECDA1TL303US262000 Rev: 07/24,V-01</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Made in India</p>  <p>10844</p>  <p>31707561085119</p> <p>12007116</p>
--	---	---

NDC 70756-288-11

Baclofen


Tablets, USP

10 mg

Rx only

100 TABLETS

Product of Poland

<p>NDC 70756-288-11</p> <p>Baclofen Tablets, USP</p> <p>10 mg</p> <p>Rx only 100 Tablets</p> 	<p>Each tablet contains Baclofen, USP 10 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECXBTLJKJKUS176800 Rev: 12/21,V-02</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Product of Poland</p>  <p>10130</p>  <p>3170756288114</p> <p>12004957</p>  <p>GTIN SN EXP LOT</p> <p>45x15 mm Unvarnished area</p>
---	--	--

NDC 70756-288-11

Baclofen

Tablets, USP

10 mg

Rx only

100 TABLETS

Made in India

<p>NDC 70756-288-11</p> <p>Baclofen Tablets, USP</p> <p>10 mg</p> <p>Rx only 100 Tablets</p> 	<p>Each tablet contains Baclofen, USP 10 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194 ECDA2TL303US262100 Rev: 07/24,V-01</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Made in India</p>  <p>10845</p>  <p>12007117</p>
---	---	--

NDC 70756-288-12

Baclofen


Tablets, USP

10 mg

Rx only

1000 Tablets

Product of Poland


<p>NDC 70756-288-12</p> <p>Baclofen Tablets, USP</p> <p>10 mg</p> <p>Rx only 1000 Tablets</p> 	<p>Each tablet contains Baclofen, USP 10 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>This package is not child resistant.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194 ECXCTLJKJKUS176900 Rev: 12/21,V-03</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Product of Poland</p>  <p>10131</p>  <p>12004980</p>
--	---	--

NDC 70756-288-12

Baclofen

Tablets, USP

10 mg
Rx only
1000 Tablets
Made in India

<p>NDC 70756-288-12</p> <p>Baclofen Tablets, USP</p> <p>10 mg</p> <p>Rx only 1000 Tablets</p> <p></p>	<p>Each tablet contains Baclofen, USP 10 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>This package is not child resistant.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECDA3TL303US262200 Rev: 07/24,V-01</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Made in India</p> <p>12007118</p>  <p>3 70756 128812 1</p>  <p>10846</p> 
--	--	--

NDC 70756-289-11
Baclofen
Tablets, USP
20 mg
Rx only
100 Tablets
Product of Poland

<p>NDC 70756-289-11</p> <p>Baclofen Tablets, USP</p> <p>20 mg</p> <p>Rx only 100 Tablets</p> <p></p>	<p>Each tablet contains Baclofen, USP 20 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECXFLLJKJKUS177100 Rev: 12/21,V-02</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Product of Poland</p> <p>12004959</p>  <p>10132</p>  <p>3 70756 128911 1</p>  <p>GTIN SN EXP LOT</p> <p>45x15 mm Unvarnished area</p>
---	--	--

NDC 70756-289-11

Baclofen
Tablets, USP
20 mg
Rx only
100 Tablets
Made in India

<p>NDC 70756-289-11</p> <p>Baclofen Tablets, USP</p> <p>20 mg</p> <p>Rx only 100 Tablets</p> <p></p>	<p>Each tablet contains Baclofen, USP 20 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECDA4TL303US262300 Rev: 07/24,V-01</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Made in India</p> <p>12007119</p>  <p>10847</p>  <p>N 3 7 0 7 5 6 1 2 8 9 1 1 1</p>
---	--	---

NDC 70756-289-51
Baclofen
Tablets, USP
20 mg
Rx only
500 Tablets
Product of Poland

<p>NDC 70756-289-51</p> <p>Baclofen Tablets, USP</p> <p>20 mg</p> <p>Rx only 500 Tablets</p> <p></p>	<p>Each tablet contains Baclofen, USP 20 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>This package is not child resistant.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECXGTLJKJKUS177200 Rev: 12/21,V-03</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Product of Poland</p> <p>12004961</p>  <p>N 3 7 0 7 5 6 1 2 8 9 5 1 1 7</p>  <p>10133</p>
---	--	---

NDC 70756-289-51

Baclofen

Tablets, USP

20 mg

Rx only

500 Tablets

Made in India

<p>NDC 70756-289-51</p> <p>Baclofen Tablets, USP</p> <p>20 mg</p> <p>Rx only 500 Tablets</p> <p>Lifestar <i>Serving Life</i></p>	<p>Each tablet contains Baclofen, USP 20 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>This package is not child resistant.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECDA5TL303US262400 Rev: 07/24,V-01</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Made in India</p> <p>12007120</p> 
--	--	---

NDC 70756-289-12

Baclofen

Tablets, USP

20 mg

Rx only

1000 Tablets

Product of Poland

<p>NDC 70756-289-12</p> <p>Baclofen Tablets, USP</p> <p>20 mg</p> <p>Rx only 1000 Tablets</p> 	<p>Each tablet contains Baclofen, USP 20 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>This package is not child resistant.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECXHTLJKJKUS177300 Rev: 12/21,V-03</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Product of Poland</p>   <p>12004962</p>
---	---	---

NDC 70756-289-12

Baclofen


Tablets, USP

20 mg

Rx only

1000 Tablets

Made in India

<p>NDC 70756-289-12</p> <p>Baclofen Tablets, USP</p> <p>20 mg</p> <p>Rx only 1000 Tablets</p> 	<p>Each tablet contains Baclofen, USP 20 mg</p> <p>USUAL DOSAGE: See package insert for full prescribing information.</p> <p>Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]</p> <p>Dispense in a well-closed container as defined in the USP, with a child-resistant closure.</p> <p>This package is not child resistant.</p> <p>Keep this and all medication out of the reach of children.</p> <p>Code No.: HP/Drugs/N-MNB/17/194</p> <p>ECDA6TL303US262500 Rev: 07/24,V-01</p>	<p>Manufactured for: Lifestar Pharma LLC 1200 MacArthur Blvd. Mahwah, NJ 07430 USA</p> <p>Made in India</p>   <p>12007121</p>
---	---	---

NDC 70756-288-01

Baclofen


Tablet, USP

10 mg

Rx only


10 tablets

Product of Poland



NDC 70756-288-01

**Baclofen
Tablet, USP
10 mg
Rx only
10 tablets
Made in India**



NDC 70756-288-99
Baclofen
Tablets, USP
10 mg per tablet
Rx only
100 Tablets
Product of Poland



NDC 70756-288-99

Baclofen


Tablets, USP

10 mg per tablet

Rx only

100 Tablets

Made in India



NDC 70756-289-01

Baclofen


Tablet, USP

20 mg


Rx only

10 Tablets

Product of Poland



Baclofen
Tablet, USP
20 mg
Rx only
10 Tablets
Made in India



NDC 70756-289-99
Baclofen
Tablets, USP
20 mg per tablet
Rx only
100 Tablets
Product of Poland



NDC 70756-289-99

Baclofen

Tablets, USP

20 mg per tablet

Rx only

100 Tablets

Made in India

Product Characteristics

Color	WHITE (white to off white)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	LS;85
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70756-085-11	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215885	01/26/2022	

BACLOFEN

baclofen tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70756-288
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)	BACLOFEN	10 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	

Product Characteristics

Color	WHITE (white to off white)	Score	2 pieces
Shape	ROUND	Size	9mm

Flavor		Imprint Code	LS;288	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70756-288-11	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2022	
2	NDC:70756-288-12	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2022	
3	NDC:70756-288-99	10 in 1 CARTON	03/25/2024	
3	NDC:70756-288-01	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA215885	01/26/2022		

BACLOFEN			
baclofen tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70756-289
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)	BACLOFEN	20 mg	
Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)			
Product Characteristics			
Color	WHITE (white to off white)	Score	2 pieces

Shape	ROUND	Size	11mm
Flavor		Imprint Code	LS;289
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70756-289-11	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2022	
2	NDC:70756-289-51	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2022	
3	NDC:70756-289-12	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/26/2022	
4	NDC:70756-289-99	10 in 1 CARTON	03/25/2024	
4	NDC:70756-289-01	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215885	01/26/2022	

Labeler - Lifestar Pharma LLC (080268943)

Registrant - Mankind Pharma Limited (915834068)

Establishment

Name	Address	ID/FEI	Business Operations
Mankind Pharma Limited		916512493	MANUFACTURE(70756-085, 70756-288, 70756-289) , ANALYSIS(70756-085, 70756-288, 70756-289) , PACK(70756-085, 70756-288, 70756-289)