

**FLU HBP MAXIMUM STRENGTH- acetaminophen, chlorpheniramine maleate,  
dextromethorphan hbr tablet, film coated**

**Walgreen Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Walgreens 44-738**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Dextromethorphan HBr 10 mg

***Purpose***

Pain reliever/fever reducer

Antihistamine

Cough suppressant

***Uses***

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - sneezing
  - headache
  - runny nose
  - cough
- temporarily reduces fever

***Warnings***

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore Throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

#### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

#### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### **When using this product**

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

#### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

#### **If pregnant or breast-feeding,**

ask a health professional before use.

#### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

#### **Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number
- protect from excessive moisture

***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

***Questions or comments?***

**1-800-426-9391**

**Principal Display Panel**

***Walgreens***

NDC 0363-7380-09

Compare to Coricidin®  
HBP Maximum Strength  
Flu active ingredients††

**MAXIMUM STRENGTH**

**Flu HBP**

**ACETAMINOPHEN /**

**PAIN RELIEVER / FEVER REDUCER**

**CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE**

**DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT**

**DECONGESTANT FREE**

- Relieves body aches, pains, headaches, runny nose, fever, cough, sore throat & sneezing
- For People with High Blood Pressure

**20**

**CAPLETS**

**ACTUAL SIZE**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

Walgreens Pharmacist Recommended

Walgreens Pharmacist Survey

††This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Flu.

50844 ORG101973809

DISTRIBUTED BY: WALGREENS CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
WALGREENS  
100% SATISFACTION  
©2020 Walgreen Co. walgreens.com

Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey  
This product is not manufactured or distributed by  
Bayer Healthcare LLC, owner of the registered  
trademark Coricidin® HBP Maximum Strength Flu.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING

NDC 0363-7380-09

Compare to Coricidin®  
HBP Maximum Strength  
Flu active ingredients††

## MAXIMUM STRENGTH Flu HBP

ACETAMINOPHEN /  
PAIN RELIEVER / FEVER REDUCER  
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE  
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DECONGESTANT FREE

- Relieves body aches, pains, headaches,  
runny nose, fever, cough, sore throat & sneezing
- For People with High Blood Pressure

20

CAPLETS

ACTUAL SIZE

KEEP OUTER PACKAGE FOR  
COMPLETE PRODUCT INFORMATION

**Drug Facts**  
Active ingredients (in each caplet)  
Acetaminophen 325 mg ..... Pain reliever/fever reducer  
Chlorpheniramine maleate 2 mg ..... Antihistamine  
Dextromethorphan HBr 10 mg ..... Cough suppressant

**Uses**  
Temporarily relieves these common cold and flu symptoms:  
■ sore throat ■ runny nose ■ cough  
■ sneezing ■ headache  
■ temporarily reduces fever

**Drug Facts (continued)**

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ blisters ■ rash ■ skin reddening  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients  
**Ask a doctor before use if you have**  
■ difficulty in urination due to enlargement of the prostate gland  
■ liver disease ■ glaucoma  
■ cough that occurs with too much phlegm (mucus)  
■ a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Drug Facts (continued)**

**When using this product**  
■ avoid alcoholic beverages  
■ marked drowsiness may occur  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ excitability may occur, especially in children  
■ use caution when driving a motor vehicle or operating machinery  
**Stop use and ask a doctor if**  
■ pain or cough gets worse or lasts more than 7 days  
■ new symptoms occur  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
**Other information**  
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

**Drug Facts (continued)**

FDCC red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Inactive ingredients** corn starch, croscopolone.

**Drug Facts (continued)**

Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ see end flap for expiration date and lot number  
■ protect from excessive moisture

**Questions or comments?** 1-800-426-9391

ORG0220

50844 ORG101973809

No Print or Varnish  
Lot and Exp. Date ONLY

ITEM 726103 W00000-0000-0

3 11917 21475 7

B-2201-738-09-H  
ORG101973809

Walgreens 44-738

## FLU HBP MAXIMUM STRENGTH

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-7380
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

CROSPVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	RED	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	44;738
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-7380-09	2 in 1 CARTON	03/19/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	03/19/2020	

**Labeler** - Walgreen Company (008965063)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	MANUFACTURE(0363-7380) , PACK(0363-7380)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(0363-7380)