

SENSICAINE™ EXTRA TOPICAL ANESTHETIC- benzocaine gel
Safco Dental Supply Co.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SensiCaine™ Extra Topical Anesthetic Gel Raspberry Flavor

Active ingredients (in each gram)

Benzocaine 200 mg

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsen, see your dentist or doctor promptly.

Do not use

- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

If pregnant or breast feeding, consult a doctor before use.

KEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions

- adults and children 2 years of age and older: apply only amount needed to the oral mucosa to prevent or relieve pain
- children under 2 years of age: do not use

Other information

- Store at 59 - 86°F (15 - 30°C).
- Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sucralose and FD&C Blue #1, FD&C Red #40, D&C Green #5, FD&C Yellow #5 as a color additive.

Questions or comments?

800-621-2178

www.safcodental.com

PRINCIPAL DISPLAY PANEL - Jar Label

SAFCO

SensiCaine™ Extra

Topical Anesthetic Gel

Contains 20% Benzocaine

Raspberry Flavor

For Professional Use Only

1 oz. (34 g)

Reorder #102-0467

NDC 67239-0217-1

Distributed by:
Safco Dental Supply Co.
Buffalo Grove, IL 60089 ☎ 800-621-2178
www.safcodental.com Made in USA

safco

SensiCaine™ Extra
Topical Anesthetic Gel

Contains 20% Benzocaine

Raspberry Flavor

1 oz. (29.6mL) For professional use only

INGREDIENTS: Each gram of 20% benzocaine gel contains between 180-220 mg benzocaine U.S.P., in a water soluble base of PEG 3350 U.S.P., PEG 400 U.S.P., flavor, sucralose, and FD&C Blue #1.

1800016(01)

Lift Here ▲

Drug Facts

Active ingredients (in each gram)

Benzocaine 200 mg

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia

warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.

Sore throat warning: If sore

fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsen, see your dentist or doctor promptly.

Do not use

- for teething
- in children under 2 years of age

When using this product

Avoid contact with eyes. If contact occurs, flush with water.

If pregnant or breast feeding, consult a doctor before use.

KEEP OUT OF REACH OF

CHILDREN. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions

- adults and children 2 years of age and older: apply only amount needed to the oral mucosa to prevent or relieve pain
- children under 2 years of age: do not use

Other information

- Store at 59 - 86° F (15 - 30° C).
- Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sucralose. May contain FD&C Blue #1, FD&C Red #40, D&C Green #5, FD&C Yellow #5 as a color additive.

Questions or comments?

800-621-2178 •

www.safcodental.com

Sore throat warning. If sore throat is severe, persists for more than 2 days, is accompanied or followed by



SENSICAINE TM EXTRA TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67239-0217
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Benzocaine (UNII: U3RSY48JW5) (Benzocaine - UNII:U3RSY48JW5)	Benzocaine	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
Polyethylene Glycol 3350 (UNII: G2M7P15E5P)	
Polyethylene Glycol 400 (UNII: B697894SGQ)	
Sucralose (UNII: 96K6UQ3ZD4)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	RASPBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67239-0217-1	29 g in 1 JAR; Type 0: Not a Combination Product	04/30/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part356	04/30/2017	

Labeler - Safco Dental Supply Co. (020417380)

Registrant - Safco Dental Supply Co. (020417380)

Revised: 11/2018

Safco Dental Supply Co.