# DAYTIME NIGHTTIME SEVERE COLD AND FLU RELIEF- acetaminophen dextromethorphan hbr guaifenesin phenylephrine hci doxylaminesucinate Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)

-----

#### **Drug Facts**

## Active ingredients (in each 15 mL) DAYTIME

Acetaminophen 325 mg Dextromethorphan HBr 10 mg Guaifenesin 200 mg

Phenylephrine HCl 5 mg

## Active ingredients for (in each 30 mL) NIGHTTIME

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine Succinate 12.5 mg
Phenylephrine HCl 10 mg

#### **Purposes for Day Time**

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

## **Purpose for Night Time**

Pain reliever/fever reducer Cough suppressant Antihistamine Nasal decongestant

#### **Uses**

#### **DAYTIME**

- temporarily relieves common cold and flu symptoms
  - nasal congestion
  - sinus congestion and pressure
  - minor aches and pains
  - headache
  - fever
  - sore throat
  - reduces swelling of nasal passages
  - cough due to minor throat and bronchial irritation
  - temporarily restores freer breathing through the nose
  - promotes nasal and/or sinus drainage
  - help loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

## **NIGHTTIME**

- temporarily relieves these common cold/flu symptoms
  - nasal congestion

- sore throat
- headache
- sinus congestion and pressure
- minor aches and pains
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- temporarily reduces
  - o fever
  - · cough to help you sleep
  - swelling of nasal passages
  - temporarily restores freer breathing through the nose
  - promotes nasal and/or sinus drainage

### **Warnings**

#### **DAYTIME**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **NIGHTTIME**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

if a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

#### **DAYTIME NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

## Ask a doctor before use if you have

#### **DAYTIME**

- liver disease
- diabetes

- heart disease
- high blood pressure
- thyroid disease
- a sodium-restricted diet
- · trouble urinating due to an enlarged prostate gland
- · persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

#### **NIGHTTIME**

- liver disease
- heart disease
- thyroid disease
- high blood pressure
- diabetes
- glaucoma
- a sodium-restricted diet
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- · trouble urinating due to enlarged prostate gland

## Ask a doctor or pharmacist before use if you are

#### **DAYTIME**

taking the blood thinning drug warfarin.

#### **NIGHTTIME**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

## When using this product

#### **DAYTIME**

#### do not take more than directed

## **NIGHTTIME**

- · do not take more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

#### Stop use and ask a doctor if

#### **DAYTIME**

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 5 days(children) or 7 days (adult)
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

#### **NIGHTTIME**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash, or headache that lasts.

These could be signs of a serious condition.

#### If pregnant or breast-feeding,

ask a health professional before use.

#### Keep out of reach of children.

#### **DAYTIME NIGHTTIME**

**Overdose warning:** Taking more than the recommended dose (overdose) may couse liver damage. In case of overdose, get medical help or contact a Poison Control Center(1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

## **DAYTIME**

- do only as directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL= milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

#### **NIGHTTIME**

- do only as directed (see Overdose warning)
- Do not exceed 4 doses per 24
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 4 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label or ensure correct dosing

#### Other information

#### **DAYTIME**

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°). Do not refrigerate

#### **NIGHTTIME**

- each 30 mL contains: sodium 64 mg
- store between 20-25°C (68-77°F). Do not refrigerate

## **Inactive ingredients**

## **Day Time**

citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xantham gum

## **Night Time**

anhydrous citric acid, FD&C blue #1, Fd&C red #40, flavor, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, trisodium citrate dehydrate, sorbitol, sucralose, xanthan gum

## Questions or comments?

Call 1-888-309-9030

## **Principal Display Panel**

## Compare to the active ingredients of VICKS® DAYQUIL® and NYQUIL® Severe Cold & Flu\* DAYTIME

Severe Cold & Flu Relief

Acetaminophen 325 mg Pain Reliever / Fever Reducer

Dextromethorphan HBr 10 mg Cough Suppressant

Guaifenesin 200 mg Expectorant

Phenylephrine HCI 5 mg Nasal Decongestant

- Headache, fever, sore throat, minor aches & pains
- Nasal/sinus congestion & sinus pressure
- Cough
- Chest congestion

For ages 6 years and over

- Max strength
- Alcohol free
- Non-drowsy

#### **NIGHTTIME**

Severe Cold & Flu Relief

Acetaminophen 650 mg Pain Reliever / Fever Reducer

Dextromethorphan HBr 20 mg Cough Suppressant

Doxylamine Succinate 12.5 mg Antihistamine

Phenylephrine HCI 10 mg Nasal Decongestant

- Headache, fever, sore throat, minor aches & pains
- Nasal/sinus congestion & sinus pressure
- Sneezing
- · Runny nose
- Cough

For ages 12 years and over

- Max strength
- · alcohol free

FL OZ (mL)

Berry Flavor

when using other Daytime or Nighttime product, carefully read each label to ensure correct dosing

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil® and NyQuil® are registered trademarks of The Procter & Gamble Company.

## TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE,

GOODLETTSVILLE, TN 37072

## **Product Label**

## DOLLAR GENERAL HEALTH Day Time Night Time Severe Cold & Flu Relief

## **DAYTIME NIGHTTIME SEVERE COLD AND FLU RELIEF**

acetaminophen dextromethorphan hbr guaifenesin phenylephrine hci doxylaminesucinate kit

## **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:55910-931

## **Packaging**

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:55910-931-16	1 in 1 KIT: Type 0: Not a Combination Product	03/31/2018	

## **Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	237 mL
Part 2	1 BOTTLE	237 mL

## Part 1 of 2

## SEVERE COLD AND FLU RELIEF DAYTIME

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

## **Product Information**

Item Code (Source)	NDC:55910-515
Route of Administration	ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg in 15 mL	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL	

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
GLYCERIN (UNII: PDC6A3C0OX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Pā	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	03/31/2018		

## Part 2 of 2

## **SEVERE COLD AND FLU NIGHTTIME**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hci liquid

Product Information	
Item Code (Source)	NDC:55910-716
Route of Administration	ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
GLYCERIN (UNII: PDC6A3C0OX)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
WATER (UNII: 059QF0KO0R)			
SODIUM CHLORIDE (UNII: 451W47IQ8X)			
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SORBITOL (UNII: 506T60A25R)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
XANTHAN GUM (UNII: TTV12P4NEE)			

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging							
#	Item Code	Package Description		Marketing Start Date		Marketing End Date	
1		237 mL in 1 BOTTLE; Type 0: Not a Combination Product					
Marketing Information							
Marketing Categ		egory	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug		ıg	M012		03/31/2018		
Marketing Information							
Marketing Category		egory	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug		ıg	M012		03/31/2018		

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 10/2023 Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)