BENZO-JEL- benzocaine gel Henry Schein Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200 mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep Out of Reach of Children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59°-86°F (15-30°C). Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

1-800-472-4346

Distributed by (in US only): HENRY SCHEIN INC. 135 DURYEA ROAD MELVILLE, NY 11747 USA

For Professional Use Only

Made in USA



mouth symptoms do not improve directed by a physician. If sore

if swelling, rash or fever develops, redness persists or worsens; or in 7 days; irritation, pain, or see your physician promptly for teething

contact occurs, flush with water. When using this product Avoid contact with eyes. If

in children under 2 years of age

Do not exceed recommended Poison Control Center right away. dosage. If more than used for pain is accidentally swallowed, get medical help or contact a

If pregnant or breast feeding (eep out of reach of children. ask a physician before use.

 Apply only amount needed to the oral mucosa to prevent or Directions

 children under 2 years of age: relieve pain. do not use

Other information

Store at 59° - 86°F (15 - 30°C). Protect from freezing.

yellow #5, (tartrazine), yellow #6, contain blue #1, green #3, green flavoring, PEG 3350, PEG 400, sodium saccharin, water. May #5, red #3, red #28, red #40, **Inactive** ingredients

Questions or comments? as a color additive.

99093451 Rev 2018/06 -800-472-4346

Oral Anesthetic

Purpose

Benzocaine 200 mg (in each g)

Active ingredients

Drug Facts

For oral mucosal use only, as emporary relief of pain due directed by dentist. For the

o minor dental procedures. Methemoglobinemia Warnings

may cause methemoglobinemia, a serious condition that must be attention if you or a child in your occur even if you have used this reduces the amount of oxygen warning: Use of this product product before. Stop use and carried in the blood. This can treated promptly because it seek immediate medical

pale, gray or blue colored skin headache (cyanosis)

care develops:

dizziness or lightheadedness shortness of breath · rapid heart rate

fatique or lack of energy

Allergy Alert: Do not use on allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine' patients with a history of

Do not use anesthetics.

for more than 7 days unless

BENZO-JEL

benzocaine gel

Product Information

HUMAN OTC DRUG NDC:0404-0033 **Product Type** Item Code (Source)

DENTAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZ OCAINE** 200 mg in 1 g

Inactive Ingredients

Ingredient Name Strength POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)

SACCHARIN SODIUM (UNII: SB8ZUX40TY)

POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)

WATER (UNII: 059QF0KO0R)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor	STRAWBERRY	Imprint Code		
Contains				

ı	P	ackaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0404-0033- 30	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018	

Marketing Information				
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
part356	05/01/2016			
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date		

Labeler - Henry Schein Inc. (012430880)

Revised: 1/2022 Henry Schein Inc.