

# **ASPIRIN LOW DOSE CHEWABLE ORANGE- aspirin tablet, chewable**

## **MAJOR PHARMACEUTICALS**

-----

### **Drug Facts**

#### **Active ingredient (in each tablet)**

Aspirin 81mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever

#### **Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for Chewable Aspirin 81 mg

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever get worse or lasts more than 3 days
  - redness or swelling is present
  - ringing in the ears or loss of hearing occurs
  - any new symptoms occur

These could be signs of a serious condition

**If pregnant or breast-feeding**

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. (800-222-1222).

**Directions**

- drink a full glass of water with each dose

---

adults and children 12 years and over	take 4 to 8 tablets every 4 hours unless directed by a doctor	not exceed 48 tablets in 24 hours
children under 12 years	consult a doctor	

---

## Other information

- store at room temperature (15<sup>0</sup>-30<sup>0</sup>C)
- **Tamper Evident:** do not use if safety seal under cap is broken or missing

## Inactive ingredients

corn starch, dextrose monohydrate, FD&C yellow #6, orange flavor, silicon dioxide, sodium saccharine, stearic acid

## Questions or comments?

**(866) 562-2756** (Mon-Fri 8 AM to 4 PM EST)

Compare to the active ingredient in Bayer® Chewable Aspirin\*\*

NDC 0904-6794-80

ASPIRIN

LOW DOSE

CHEWABLE

ORANGE

PAIN RELIEVER

(NSAID)

81mg

1000 TABLETS

Compare to the active ingredient in Bayer® Chewable Aspirin\*\*

NDC 0904-6794-80

ASPIRIN  
LOW DOSE  
CHEWABLE  
ORANGE  
PAIN RELIEVER  
(NSAID)

81mg

1000 Tablets

MAJOR®

<b>Drug Facts</b>	<p><b>Active ingredient (in each tablet)</b> Aspirin 81 mg (NSAID) <span style="font-size: x-x-small;">Pain reliever</span></p> <p><b>Nonsteroidal anti-inflammatory drug</b></p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>■ for the temporary relief of minor aches and pains or as recommended by your doctor</li> <li>■ ask your doctor about other uses for Chewable Aspirin 81 mg</li> </ul> <p><b>Warnings</b></p> <p><b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in mental status, persistent vomiting, or other symptoms occur, these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing aspirin, ibuprofen, or acetaminophen (NSAIDs, aspirin, blood thinners, or other) ■ take more than 3 alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p><b>Do not use</b> ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor</p> <p><b>Ask a doctor before use if</b> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease ■ you have asthma ■ you are taking a diuretic</p> <p><b>Ask a doctor or pharmacist before use if you are</b> taking a prescription drug for ■ gout ■ diabetes ■ arthritis</p> <p><b>Stop use and ask a doctor if</b> ■ an allergic reaction occurs. Seek medical help right away ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ ringing in the ears or loss of hearing</p> <p><b>These could be signs of a serious condition:</b></p> <p>■ pregnant or breast-feeding, ask a health professional before use, <b>it is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during or after birth</b> ■ you are taking aspirin for a headache, get medical help or contact a Poison Control Center right away (800-222-1222).</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ drink a full glass of water with each dose</li> <li>adults and children 12 years and over take 4 to 8 tablets every 4 hours not exceed 48 tablets in 24 hours unless directed by a doctor</li> <li>children under 12 years consult a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store at room temperature (15°-30°C) ■ <b>Tamper Evident:</b> do not use if safety seal under cap is broken or missing</li> </ul> <p><b>Inactive ingredients</b> corn starch, dextrose monohydrate, FD&amp;C yellow #6, orange flavor, silicon dioxide, sodium saccharine, stearic acid</p> <p><b>Questions or comments? (866) 562-2756</b> (Mon-Fri 8 AM to 4 PM EST)</p>
-------------------	---

\*\*This product is not manufactured or distributed by Bayer®, owner of the registered trademark Bayer® Chewable Aspirin.

Distributed by:  
**MAJOR PHARMACEUTICALS**  
Indianapolis, IN 46208  
www.majorpharmaceuticals.com  
(800) 615-2471

Rev. 12/23 Re-order No. 701003 M-111



3 0904679480 6

SPOT UV

## ASPIRIN LOW DOSE CHEWABLE ORANGE

aspirin tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0904-6794
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	PH;034
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6794-89	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/27/2018	
2	NDC:0904-6794-80	1000 in 1 PACKAGE; Type 0: Not a Combination Product	09/27/2018	
3	NDC:0904-6794-30	300 in 1 CARTON	02/13/2023	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/27/2018	

**Labeler** - MAJOR PHARMACEUTICALS (191427277)**Registrant** - Pharbest Pharmaceuticals, Inc. (557054835)**Establishment**

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Pharbest Pharmaceuticals, Inc.

557054835

manufacture(0904-6794)

Revised: 12/2025

MAJOR PHARMACEUTICALS