

**BACKAID MAX- acetaminophen, pamabrom tablet, film coated**  
**Kobayashi Healthcare International, Inc.**

-----

**BackAid**

**Active ingredients (in each caplet)**

Acetaminophen, 500 mg

Pamabrom, 25 mg

**Purpose**

Analgesic (pain reliever)

Diuretic

**Uses**

For the temporary relief of:

- minor aches, pains and related discomforts due to muscle strain, spasms or overexertion including those affecting the back, legs and joints
- pressure-caused discomforts due to periodic excess water retention

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

**Ask a doctor before use if you** have liver disease.

**Ask a doctor or pharmacist before use if you** are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a poison control center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- Read all package directions and warnings before use. Use only as directed.
- Adults: Two (2) caplets with a full glass of water every 6 hours while symptoms persist, not to exceed 3 doses (6 caplets) in 24 hours, or as directed by a doctor.
- Do not exceed recommended dosage.
- For use by normally healthy adults only.
- Persons under 18 years of age should use only as directed by a doctor.

**Other information**

- Magnesium content: 5 mg/caplet
- **\*\*CONTENTS SEALED:** Each BACKAID MAX golden-colored, oval shaped caplet bears the identifying mark "ALVA" and is sealed in a clear plastic blister with a paper/foil backing. Do not use if seal appears broken or if product contents do not match product description.
- You may report serious side effects to the phone number provided under *Questions?* below.

**Inactive ingredients** Calcium sulfate, croscarmellose sodium, D&C yellow #10 Lake, FD&C yellow #6 Lake, hypromellose, magnesium oxide, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, polyvinylpyrrolidone, potassium gluconate, sodium lauryl sulfate, sodium starch glycolate, sorbitol, starch, stearic acid, talc, titanium dioxide.

**Questions? 1-800-792-2582.**

**Package insert**

Aspirin Free

Backache Relief

**BACKAID Max**

1000 mg Strong per 2 Caplet Dose

## Maximum Strength 6 Hour Formula

**Active ingredients:** Each caplet contains: acetaminophen 500 mg (analgesic/pain reliever) and pamabrom 25 mg (diuretic).

**Uses:** For the temporary relief of: • minor aches, pains and related discomforts due to muscle strain, spasms or overexertion including those affecting the back, legs and joints. • pressure-caused discomforts due to periodic excess water retention.

### Warnings

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** • with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • for more than 10 days for pain unless directed by a doctor • for more than 3 days for fever unless directed by a doctor.

**Ask a doctor before use if you** have liver disease.

**Ask a doctor or pharmacist before use if** you are taking the blood thinning drug warfarin.

### Stop use and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a poison control center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions:

- Read all package directions and warnings before use. Use only as directed.
- Adults: Two (2) caplets with a full glass of water every 6 hours while symptoms persist, not to exceed 3 doses (6 caplets) in 24 hours, or as directed by a doctor.
- Do not exceed recommended dosage.
- For use by normally healthy adults only.
- Persons under 18 years of age should use only as directed by a doctor.

### Other information:

Magnesium content: 5 mg/caplet

**\*\*CONTENTS SEALED:** Each BACKAID MAX golden-colored, oval shaped caplet bears the identifying mark "ALVA" and is sealed in a clear plastic blister with a paper/foil backing. Do not use if seal appears broken or if product contents do not match product

description. • You may report serious side effects to the phone number provided under *Questions?* below.

**Inactive ingredients** Calcium sulfate, croscarmellose sodium, D&C yellow #10 Lake, FD&C yellow #6 Lake, hypromellose, magnesium oxide, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, polyvinylpyrrolidone, potassium gluconate, sodium lauryl sulfate, sodium starch glycolate, sorbitol, starch, stearic acid, talc and titanium dioxide.

**Questions? 1-800-792-2582**

53-38RC,  
0424

- Tough on Pain,  
Easy on the Stomach™
- **ASPIRIN FREE**

*Vea las instrucciones  
en español adentro*

**Maximum Strength  
BACKAID®**

ASPIRIN FREE  
1000 MG STRONG  
PER 2 CAPLET DOSE  
Contains no  
gluten ingredients

**Maximum Strength  
BACKAID®**



**Eases Pain of  
Inflammation**  
**Relieves Backache  
and Sciatica Pain**


**6** Hour Formula

**28** CAPLETS

**Acetaminophen + Pamabrom**

Prod No.: 53028  
**Contents Sealed\*\***

Lot No.:  
Exp.:




KOBAYASHI Consumer Products, LLC  
P.O. Box 1191, Dalton, GA 30722  
© 1988-2024 ALVA  
® Reg. U.S. Patent and Trademark Office



Recyclable

[www.BACKAID.com](http://www.BACKAID.com)



0 72959 53028 0

**Stop use and ask a doctor if** ■ symptoms do not improve ■ new symptoms occur ■ pain or fever persists or gets worse

**Keep out of reach of children.** In case of overdose, get medical help or contact a poison control center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ Read all package directions and warnings before use. Use only as directed.

■ Adults: Two (2) caplets with a full glass of water every 6 hours while symptoms persist, not to exceed 3 doses (6 caplets) in 24 hours, or as directed by a doctor. ■ Do not exceed recommended dosage. ■ For use by normally healthy adults only. ■ Persons under 18 years of age should use only as directed by a doctor.

**Other information** ■ Magnesium content 5 mg/caplet ■ **\*\*CONTENTS SEALED.** Each BACKAID golden colored, oval shaped caplet bears the identifying mark "ALVA" and is sealed in a clear plastic blister with a paper/foil backing. Do not use if seal appears broken or if product contents do not match product description. ■ You may report serious side effects to the phone number provided under *Questions?* below.

**Inactive ingredients** Calcium sulfate, croscarmellose sodium, D&C yellow #10 Lake, FD&C yellow #6 Lake, hypromellose, magnesium oxide, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, polyvinylpyrrolidone, potassium gluconate, sodium lauryl sulfate, sodium starch glycolate, sorbitol, starch, stearic acid, titanium dioxide.

**Questions? 1-800-792-2582**


PMS 3005

PMS 2126

PMS 186

PMS 108

BLACK



## BACKAID MAX

acetaminophen, pamabrom tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52389-153
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
PAMABROM (UNII: UA8U0KJM72) (BROMOTHEOPHYLLINE - UNII:FZ G87K1MQ6)	PAMABROM	25 mg

### Inactive Ingredients

Ingredient Name	Strength
CALCIUM SULFATE DIHYDRATE (UNII: 4846Q921YM)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
TALC (UNII: 7SEV7J4R1U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POTASSIUM GLUCONATE (UNII: 12H3K5QKN9)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SORBITOL (UNII: 506T60A25R)	
STEARIC ACID (UNII: 4ELV7Z 65AP)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
POVIDONE (UNII: FZ989GH94E)				
Product Characteristics				
Color	orange (Goldenrod)		Score	no score
Shape	OVAL		Size	15mm
Flavor			Imprint Code	ALVA
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52389-153-28	2 in 1 CARTON	03/01/2011	
1	NDC:52389-153-01	14 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:52389-153-14	1 in 1 CARTON	06/01/2011	
2	NDC:52389-153-01	14 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:52389-153-38	1 in 1 CARTON	09/01/2010	
3	NDC:52389-153-03	38 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:52389-153-46	1 in 1 CARTON	04/29/2016	
4		46 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:52389-153-35	5 in 1 CARTON	06/01/2011	
5		7 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	02/01/1992	

**Labeler** - Kobayashi Healthcare International, Inc. (156391729)

**Registrant** - Kobayashi America Manufacturing, LLC (079852150)