WELBY INFANTS ACETAMINOPHEN- acetaminophen suspension Aldi Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Aldi Inc. Infants' Acetaminophen 160 mg per 5mL Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart			
Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	

^{*} or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

- store at 20-25°C (68-77°F)
- · do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Infants' Tylenol® Oral Suspension

active ingredient

See new warnings

For Ages 2 to 3 Years

Infants'

Pain & Fever

Acetaminophen 160mg per 5mL

Oral Suspension

Pain Reliever & Fever Reducer

Alcohol Free

Aspirin Free

Ibuprofen Free

Dye Free

Use only with enclosed syringe.

See side panel for more information.

Grape flavor

2 FL OZ (60 mL)





Please see dosing chart or ask a doctor for your child's correct dose

STEP 1. See *Directions* for the correct dose for your child.

STEP 3. Push air out of syringe. Insert syringe tip into bottle opening.



STEP 4. Flip bottle upside down.
Pull yellow part of syringe to
the first dose line and then push product back into bottle

STEP 5. Pull yellow part of syringe until it reaches and stays at the correct dose.



STEP 6. Dispense liquid slowly into child's mouth



Drug Facts

Active ingredient (in each 5 mL)

Uses temporarily: ■ reduces fever
■ relieves minor aches and pains due to: ■ the common cold

■ flu ■ headache ■ sore throat

■ toothache

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash,

nausea, or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

sup use and ask a doctor if we had 5 days
■ fever gets worse or lasts more than 3 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ ™ erdenses or swelling is present.
These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed (see overdose warning)
 ■ shake well before using ■ mL = milliliter
- find right dose on chart. If possible, use weight to dose;

- into right cose on clark. If possible, use weight to dose, otherwise, use age.
 push air cut of syringe. Insert syringe tip into bottle opening.
 flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.

Drug Facts (continued)

- pull yellow part of syringe until it reaches and stays at the correct dose
 dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while
- symptoms last
 do not give more than 5 times in 24 hours
 replace cap tightly to maintain

child resistance Dosing Chart

ı	Weight (lb)		Dose (mL)*
ı	under 24	under 2 years	ask a doctor
l	24-35	2-3 years	5 mL

or as directed by a doctor Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing

Other information

- store at 20-25°C (68-77°F)
 do not use if printed neckband is
- broken or missing

Inactive ingredients anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, navor, glycenn, nydrokyerniy cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylen glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

distributor of Infants' Tylenol® Oral Suspension

DISTRIBUTED BY ALDI INC BATAVIA, IL 60510







LOT NO.

FXP.

: 76636 3D C3

WELBY INFANTS ACETAMINOPHEN

acetaminophen suspension

Product Information

HUMAN OTC DRUG NDC:64024-766 Product Type Item Code (Source)

ORAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0 DDB505)	
CARRAGEENAN (UNII: 5C69 YCD2YJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
CELLULO SE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679 OBS 311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SO DIUM PHO SPHATE, TRIBASIC (UNII: A752Q30 A6 X)	

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

ı	Packaging				
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
l	1 NDC:64024-766-16	1 in 1 CARTON	12/16/2016		
l	1	60 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/16/2016	

Labeler - Aldi Inc (944259522)

Revised: 12/2018 Aldi Inc