SIGNATURE CARE DAYTIME SEVERE COLD- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder, for solution Safeway

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Better Living Brands LLC Severe Cold Drug Facts

Active ingredients (in each packet)

Acetaminophen 500 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer Cough suppressant Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- · thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

| Age | Dose |
|--|------------|
| adults and children 12 years of age and over | one packet |
| children under 12 years of age | do not use |

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylketonurics: contains phenylalanine 22 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-888-723-3929

Package/Label Principal Display Panel

Compare to Theraflu® Multi-Symptom Severe Cold active ingredients

Quality Guaranteed

Daytime Multi-Symptom Severe Cold

_ _ _

ACETAMINOPHEN

Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr

Cough Suppressant

PHENYLEPHRINE HCI

Nasal Decongestant

GREEN TEA & HONEY LEMON FLAVORS

Nasal Congestion

Cough

Body Ache

Sore Throat Pain

Headache

Fever

6 PACKETS













Severe Cold

MADE IN MEXICO

DISTRIBUTED BY: BETTER LIVING BRANDS LLC P.O. BOX 99, PLEASANTON, CA 94566-0009 1-888-723-3929 www.betterlivingbrandsl_LC.com

> OUR PROMISE GUARANTEED



Drug Facts Active ingredients (in each packet) Purposes Dextromethorphan HBr 20 mg Phenylephrine HCl 10 mg

- Uses
 temporarily relieves these symptoms due to a cold:
 minor aches and pains minor sore throat pain
 headache masal and sirus congest
 cough due to minor throat and bronchial irritation
 temporarily reduces fever

Warnings Liver warning. This product contains acetaminophen. Severe liver damage may occur if you take in more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen in 3 or more abholici drinks every day while using this product. Alteryy alert. Acetaminophen may cause severe skin reactions. Symptoms may include: in skin reddering in blotters in radio. If you have been product and a seek medical help right away.

- In sain reaction occurs, stop use and seek medical help right away.

 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or wonlting, consult a doctor promptly.

- Do not use "in a child under 12 years of age
 If you have ever had an alterior exaction to this product or any of its ingredients
 If you have ever had an alterior reaction to this product or any of its ingredients
 with any other flug containing academinophen igneraction or renorpescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 If you are not whiting a prescription mornoamine coldses inhibitor (MAOI) (cortain drugs for degression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI frug. If you do not how if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

LOT NO. EXP : 8Y491 LJ CL

- Ask a doctor before usef fyou have been being to grow the first disease. If the first disease is heart disease is heart disease is heart disease is heart disease is the first disease is heart disease is disease is heart disease is heart disease is disease is heart disease is disease is
- a sodium-restricted diet
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

- Stop use and ask a doctor if Innervounness, dizziness, or eleoplessness occurs

 Intercepts worse or lasts more than 3 days

 Interduses or swelling is present. In thew ymptoms occur

 Inpain, ough or nead conjection gets worse or lasts more than 7 days

 Incupil come back or occurs with nark or headdard that lasts. These could be signs of a

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not use more than directed (see overdose warning) take every 4 hours, while symptoms persist.
 Do not take more than 6 packets in 24 hours
 the directed by a doctor.

| uniess un octeu by a doctor. | | |
|---|------------|--|
| Age | Dose | |
| adults and children 12 years of age and over | one packet | |
| children under 12 years of age | do not use | |

- in dissolve contents of one packet into 8 oz. hot water, sip while hot Consume entire drink within 10-15 minutes.

 If using a microwave, add contents of one packet by 8 oz. of cool water, stip ribriskly before and after heating. Do not overheat.

Other information

- sodium 25 mg
 phenylketonurios: contains phenylalanine
 22 mg per packet
 store at 20-25°C (68-77°F). Protect product from
 heat and moisture.

Inactive Ingredients acesulfame potas anhydrous citric acid, aspartame, colloidal silic dioxide, D&C yellow #10, FD&C blue #1, FD&C ed #40, flavors, mallodextrin, pregelatinized s sodium citrate, sucrose, tribasic calcium phosp

Questions or comments?

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® Multi-Symptom Severe Cold.



www.StopMedicineAbuse.org







Scan here for more



SIGNATURE CARE DAYTIME SEVERE COLD

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder, for solution

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:21130-937

Route of Administration ORAL

Active Ingredient/Active Moiety Basis of Strength Strength **Ingredient Name** ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) | ACETAMINOPHEN 500 mg **DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)** DEXTROMETHORPHAN 20 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** PHENYLEPHRINE HYDROCHLORIDE (UNII: 04|A59TNS|) (PHENYLEPHRINE -**PHENYLEPHRINE** 10 mg UNII:1WS297W6MV) **HYDROCHLORIDE**

| Inactive Ingredients | | |
|---|----------|--|
| Ingredient Name | Strength | |
| ACESULFAME POTASSIUM (UNII: 230V73Q5G9) | | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | | |
| ASPARTAME (UNII: Z0H242BBR1) | | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | | |
| SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) | | |
| SUCROSE (UNII: C151H8M554) | | |
| TRIBASIC CALCIUM PHOSPHATE (UNII: 91D9GV0Z28) | | |

| Product Characteristics | | | |
|-------------------------|---------------------------------------|--------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | HONEY (green tea) , LEMON (green tea) | Imprint Code | |
| Contains | | | |

| ı | Packaging | | | |
|---|----------------------|--|-------------------------|-----------------------|
| 4 | # Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:21130-937- 91 | 6 in 1 CARTON; Type 0: Not a Combination Product | 05/17/2019 | |

| Marketing Information | | | |
|-----------------------|---|-------------------------|-----------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC monograph final | part341 | 05/17/2019 | |
| | | | |

Labeler - Safeway (009137209)

Revised: 11/2022 Safeway