

**PANCOLD A- acetaminophen, guaifenesin, pentoxyverine citrate, phenylephrine hydrochloride, chlorpheniramine maleate, caffeine anhydrous liquid**  
**DONGWHA PHARM. CO. LTD.**

-----

**PANCOLD A ORAL SOLUTION**

**ACTIVE INGREDIENTS**

**Active ingredients (in each 30 mL bottle)**

- Acetaminophen 300 mg
- Guaifenesin 80 mg
- Pentoxyverine citrate 15 mg
- Phenylephrine Hydrochloride 10 mg
- Chlorpheniramine maleate 2.5 mg
- Caffeine 30 mg

**PURPOSE**

Pain Reliever-Fever Reducer, Expectorant, Antitussive, Nasal Decongestant, Antihistamine, Pain reliever aid

**INDICATIONS & USAGE**

**Uses** temporarily relieves these common cold symptoms:

- runny nose & sneezing
- nasal congestion
- sore throat
- cough
- phlegm
- chills
- fever
- headache
- joint pain
- muscle pain

**WARNINGS SECTION**

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more glasses of alcohol everyday while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

- **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **DO NOT USE**

### **Do not use**

- if you ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product.
- If you are taking expectorants, other cold drugs, antipyretic-analgesics, sedatives, oral drugs containing antihistamines (oral drugs for rhinitis, motion sickness drugs, and allergy drugs)

## **ASK A DOCTOR**

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)
- glaucoma (eye pain, blurred vision)
- weak body
- high fever

## **STOP USE**

### **Stop use and ask doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or last more than 7 days
- fever gets worse or lasts more than 3 days and redness or swelling is present
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

## **PREGNANCY OR BREAST FEEDING**

**If pregnant or breast-feeding, ask a health professional before use.**

## **KEEP OUT OF REACH OF CHILDREN**

**Keep out of reach of children.**

## **OVERDOSAGE**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptom.

## **DOSAGE & ADMINISTRATION**

### **Directions**

- do not take more than directed (see overdose warning)

---

Adults: Take 1 bottle (30mL) three times a day, 30 minutes after meal.  
Children and infants: Do not administer. Ask a doctor.

---

## **STORAGE AND HANDLING**

### **Other nformation**

- store at room temperature between 15°-25°C (59°-77°F) and avoid direct sunlight

## **INACTIVE INGREDIENTS**

- **Inactive Ingredients:** : alcohol, citric acid monohydrate, edetate disodium, fructooligosaccharide, lemon flavor, orange flavor, propylene glycol, sodium citrate, sucralose, water

**Drug Facts**

**Active ingredients (in each 30 mL bottle)**

- Acetaminophen 300 mg
- Chlorpheniramine maleate 2.5 mg
- Caffeine 30 mg
- Guafenesin 15 mg
- Phenylephrine hydrochloride 10 mg
- Pseudoephedrine sulfate 15 mg
- Pentoxifyverine citrate 15 mg

**Purpose**

Relieves cold symptoms

**Warnings**

Do not use if you have had an allergic reaction to the product or any of its ingredients. Do not use if you are taking a prescription drug containing MAOI. Do not use if you are taking a prescription drug containing MAOI. Do not use if you are taking a prescription drug containing MAOI.

**Directions**

Adults: Take 1 bottle (30mL) three times a day, 30 minutes after meal.

Children and Infants: Do not administer. Ask a doctor.

**Fast Relief Cold Medicine**

**판콜에이 A**

**PANCOLD**

**나쁜약**

**Fast Relief Cold Medicine**

**판콜에이 A**

**PANCOLD**

**나쁜약**

**Runny Nose**

**Nasal Congestion**

**Cough**

**Sneezing**

**Phlegm**

**Sore Throat**

**Chills**

**Fever**

**Headache**

**Muscle pain**

**1 FL OZ (30mL) X 3**

**DO NOT SELL SEPARATELY**

**Fast Relief Cold Medicine**

**판콜에이 A**

**PANCOLD**

**나쁜약**

**Runny Nose**

**Cough**

**Sore throat**

**Body aches**

**1 FL OZ (30mL) X 3**

**DO NOT SELL SEPARATELY**

## PANCOLD A

acetaminophen, guaifenesin, pentoxifyverine citrate, phenylephrine hydrochloride, chlorpheniramine maleate, caffeine anhydrous liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:51352-189
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	80 mg in 30 mL
<b>PENTOXIFYVERINE CITRATE</b> (UNII: 4SH0MFJ5HJ) (PENTOXIFYVERINE - UNII:32C726X12W)	PENTOXIFYVERINE CITRATE	15 mg in 30 mL
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	300 mg in 30 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2.5 mg in 30 mL
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	30 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	32.4 mg in 30 mL
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	10 mg in 30 mL

<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	29 mg in 30 mL
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	3 mg in 30 mL
<b>ALCOHOL</b> (UNII: 3K9958V90M)	0.5 mL in 30 mL
<b>SACCHARUM OFFICINARUM STEM FRUCTOOLIGOSACCHARIDES</b> (UNII: 8LLD82AE3S)	7.5 g in 30 mL
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ORANGE</b> (UNII: 5EVU04N5QU)	
<b>LEMON</b> (UNII: 24RS0A988O)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	ORANGE (Light yellowish clear oral liquid with orange-lemon flavor.)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51352-189-73	3 in 1 CASE	07/01/2023	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:51352-189-75	5 in 1 CASE	12/07/2023	12/07/2023
2		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2023	

**Labeler** - DONGWHA PHARM. CO. LTD. (687745240)

**Registrant** - DONGWHA PHARM. CO. LTD. (687745240)

### Establishment

Name	Address	ID/FEI	Business Operations
DONGWHA PHARM. CO., LTD.		687745240	manufacture(51352-189)

Revised: 12/2024

DONGWHA PHARM. CO. LTD.